



Department
for Environment
Food & Rural Affairs

Pest specific plant health response plan:

Outbreaks of *Epitrix* potato flea beetles on potato
crops




Figure 1. Potato tuber showing feeding damage by the larvae of a potato flea beetle in Portugal. © Conceição Boavida Instituto Nacional de Recursos Biológicos, Portugal

We are the Department for Environment, Food and Rural Affairs. We're responsible for improving and protecting the environment, growing the green economy, sustaining thriving rural communities and supporting our world-class food, farming and fishing industries.

We work closely with our 33 agencies and arm's length bodies on our ambition to make our air purer, our water cleaner, our land greener and our food more sustainable. Our mission is to restore and enhance the environment for the next generation, and to leave the environment in a better state than we found it.



© Crown copyright 2022

This information is licensed under the Open Government Licence v3.0. To view this licence, visit www.nationalarchives.gov.uk/doc/open-government-licence/

This publication is available at

<https://planthealthportal.defra.gov.uk/pests-and-diseases/contingency-planning/>

Any enquiries regarding this document should be sent to us at:

The UK Chief Plant Health Officer
Department for Environment, Food and Rural Affairs
Room 11G32
York Biotech Campus
Sand Hutton
York
YO41 1LZ

Email: plantpestrisks@defra.gov.uk

www.gov.uk/defra

Executive summary

Background	
Regulation	GB Quarantine pest
Key Hosts (2.3)*	Potatoes
Distribution	See appendix A.
Key pathways	Seed and ware potatoes.
Industries at risk	Potato growers and packers.
Symptoms (2.5)	Shot hole of leaves, holes, tunnelling and splinter lesions on tuber
Surveillance	
Demarcated zones (5.24-5.28)	Infested zone = Infested field(s) Buffer zone = ≥ 1 km
Surveillance activities (5.26-5.28)	<ul style="list-style-type: none"> • Visual surveys including sweep net sampling, will be carried out in the infested zone and neighbouring fields. • Following this, surveys should be extended to potato fields and hosts within the buffer zone.
Response measures	
Interceptions (5.1-5.5)	<ul style="list-style-type: none"> • Consignment should be destroyed or re-exported if dead larvae or feeding damage is seen. • Tracing exercises carried out where required • UKPHINs notification to be made.
Outbreaks (5.32-5.46)	<p>This is dependent on the growth stage and outbreak situation.</p> <p>Measures could include:</p> <ul style="list-style-type: none"> • Foliar insecticide treatments • Monitoring of pest levels • Destruction of haulm if pest levels are low • Removal of volunteers and weeds • Harvesting restrictions and destruction of volunteers • Destruction of infested tubers
Key control measures	
Biological	N/A
Chemical	Insecticide and herbicide applications
Cultural	Cleaning of equipment and machinery, removal and destruction of waste, destruction of infested tubers
Declaration of eradication	
Eradication can be declared if no pest is detected during annual surveys for three years after the infested material was destroyed. This should include two years with no volunteers and a following year with only a ware or trap crop.	

*

Numbers refer to relevant points in the plan

Contents

Executive summary	3
1. Introduction and scope	5
2. Summary of threat.....	5
3. Risk Assessments.....	6
4. Actions to prevent outbreaks.....	6
5. Response.....	7
Official action to be taken following the suspicion or confirmation of <i>Epitrix</i> potato flea beetles on imported plants, including tubers and seeds.....	7
Official action to be taken following the suspicion of an <i>Epitrix</i> sp. outbreak.....	8
Confirming a new outbreak	10
Criteria for determining an outbreak.....	12
Official Action to be taken following the confirmation of an outbreak	12
Measures in subsequent seasons.....	17
6. Criteria for declaring eradication / change of policy	18
7. Evaluation and review of the contingency plan	18
8. Appendix A.....	19
Data sheet for <i>Epitrix similaris</i> , <i>Epitrix tuberis</i> , <i>Epitrix cucumeris</i> and <i>Epitrix papa</i>	19
9. Appendix B.....	34
Statutory Action.....	34
10. References	37
11. Authors and reviewers	40

1. Introduction and scope

- 1.1. This pest specific response plan has been prepared by the Defra Risk and Horizon Scanning team. It describes how the Plant Health Service for England will respond if an infestation of non-indigenous potato flea beetles of the genus *Epitrix* Foudras, and specifically the species *E. cucumeris*, *E. papa*, *E. similaris* and/or *E. tuberis* is discovered on potato (*Solanum tuberosum*).
- 1.2. The plant health authorities in Northern Ireland, Scotland, Wales and the Crown Dependencies have been consulted on this plan and will use it as the basis for the action they will take in the event of *E. cucumeris*, *E. papa*, *E. similaris* and *E. tuberis* being detected in their territories.
- 1.3. This document will be used in conjunction with the *Defra Contingency Plan for Plant Health in England* (<https://planthealthportal.defra.gov.uk/assets/uploads/Generic-Contingency-Plan-for-Plant-Health-in-England-FINAL-2.pdf>), which gives details of the teams and organisations involved in pest response in England, and their responsibilities and governance. It also describes how these teams and organisations work together in the event of an outbreak of a plant health pest.
- 1.4. The aim of this response plan is to facilitate the containment and eradication of *E. cucumeris*, *E. papa*, *E. similaris* and *E. tuberis*. For the purposes of this plan, these will be referred to as *Epitrix* potato beetles.

2. Summary of threat

- 2.1. *Epitrix cucumeris*, *E. similaris* and *E. tuberis* and *E. papa* belong to a group of six *Epitrix* species currently known to feed on potato. The first five species are all known to be of North American origin. The origin of *E. papa* is currently unknown, but it could also be of North American origin. In 2004, potato tuber damage indicative of *Epitrix* was observed in Europe for the first time, in north Portugal, near Porto. *Epitrix cucumeris* and a second *Epitrix* species were subsequently confirmed as the cause of this damage in 2008. The second species was at first identified as *E. similaris* but following a morphological investigation this species was recognised as a new species and named as *E. papa* in 2015 (Orlova-Bienkowskaja, 2015). This new species has now been reported from the Portugese counties of Alentejo (2017), Calheta (2018), Machico (2018), Ponta Do Sol (2018), Porto Moniz (2018) and São Vicente (2018). *Epitrix papa* has also been recorded in Spain, in Galicia (2010 – date of first record), Asturias (2014), and Andalucia (2015). *Epitrix cucumeris* has been reported from Portugal (Azores and Madeira) and Spain, whereas *E. similaris* and *E. tuberis* are reported as absent from Europe. All species are absent from the UK.

- 2.2. Adult *Epitrix* potato flea beetles feed on foliage and produce characteristic “shot-like” holes, while the larvae burrow underneath the surface of tubers, forming holes, serpentine tunnels and deep splinter lesions (Boavida *et al.*, 2013). Damage by larvae often leads to the rejection of tubers by potato packers in Europe and subsequent economic loss. The deeper holes and lesions can also render consignments unsuitable for processing.
- 2.3. In the UK, there have been no interceptions of live *Epitrix* potato beetles. However, there have been three interceptions of dead larvae on washed ware potatoes imported from Spain. *Epitrix* feeding damage has also been found on washed and unwashed ware potatoes originating from areas of Spain where the pest was not known to be present.

3. Risk Assessments

- 3.1. *Epitrix cucumeris*, *E. papa* and *E. tuberis* have a mitigated UK plant health risk register score of 30 (moderate risk). *Epitrix similaris* has a lower score of 10 (low risk) because of its very restricted distribution (localised areas in the United States) and very limited reporting of tuber damage. Overall scores range from 1 (very low risk) to 125 (very high risk). These scores are reviewed as and when new information becomes available (<https://planthealthportal.defra.gov.uk/pests-and-diseases/uk-plant-health-risk-register/index.cfm>).
- 3.2. Following the introduction of *Epitrix* potato flea beetles into Portugal, the European and Mediterranean Plant Protection Organization (EPPO) Working Party on Phytosanitary Regulations decided that a Pest Risk Analysis for *E. cucumeris*, *E. similaris*, *E. subcrinita* and *E. tuberis* should be carried out. This was completed in 2011, with modifications in 2012 (http://www.eppo.int/QUARANTINE/Pest_Risk_Analysis/PRA_intro.htm). It should be noted that the PRA refers to *E. similaris* rather than *E. papa* in Portugal and Spain because it was written before *E. papa* was recognised.

4. Actions to prevent outbreaks

- 4.1. *Epitrix cucumeris*, *E. papa* and *E. tuberis* are GB quarantine pests ([Schedule 1 of The Plant Health \(Phytosanitary Conditions\) \(Amendment\) \(EU Exit\) Regulations 2020](#)) and are therefore prohibited from being introduced into, or spread within GB. Further pest and host specific requirements are listed in [Schedule 7](#). They are also a GB Priority Pests meaning they are GB quarantine pests which has been assessed to have the most severe potential economic, environmental and social impacts to GB.

- 4.2. *Epitrix tuberis* is an EPPO A1 listed pest and *E. cucumeris* and *E. papa* are EPPO A2 listed pests. The A1 and A2 lists are species that EPPO should be regulated by EPPO countries, the A1 list for species that are absent from the region and the A2 list for species that are present in some countries. *Epitrix similis* was an A2 listed pest for the EPPO region but was removed from the list in 2016 following the recognition that the records of *E. similis* from Europe were erroneous and were in fact records of *E. papa*, a previously unknown species. Therefore, *E. similis* has never been recorded in the EPPO region.
- 4.3. *Epitrix cucumeris*, *E. papa* and *E. tuberis* are subject to EU Emergency measures to prevent their introduction and spread within the EU.
- 4.4. The Plant Health Service should be aware of the measures described in this plan and be trained in responding to an outbreak of *Epitrix* potato flea beetles. It is important that capabilities in detection, diagnosis, and risk management are available.

5. Response

Official action to be taken following the suspicion or confirmation of *Epitrix* potato flea beetles on imported plants, including tubers and seeds

Holding consignments at interception points, including packhouses

- 5.1. If *Epitrix* potato flea beetles are suspected by the Plant Health and Seeds Inspectorate (PHSI) to be present on a consignment moving in trade, the PHSI should hold the consignment (ideally in a sealed cold store or covered in a way as to not allow the escape of the beetle) until a diagnosis is made. Samples should be sent in by the PHSI to Fera Science Ltd., Plant Clinic, York Biotech Campus, Sand Hutton, York, YO41 1LZ (01904 462000) in a sealed bag or container, within at least two other layers of containment, which are not liable to be crushed during transit. If an infestation of *Epitrix* potato flea beetles is confirmed, the PHSI will advise the client of the action that needs to be taken by way of an official plant health notice. The consignment should be destroyed by either deep burial or re-exported to the place of origin in a sealed container (see also 5.48-5.50). Defra's Risk and Horizon Scanning team can advise on alternative methods of destruction if necessary.
- 5.2. The consignment should also be destroyed or re-exported if dead larvae and/or feeding damage are found on unwashed potatoes. See Appendix B for full details of statutory action on potato tubers arriving from Portugal and Spain.

- 5.3. Any host plants (including any tubers, which should be held) should be surveyed on the site or in the immediate vicinity in the summer and again in the following year for signs of pest presence (see 5.15 -5.18). If the site is in a potato growing area, the two potato fields closest to the site and a field on the prevailing wind side should also be surveyed. Waste disposal processes and areas should likewise be inspected.
- 5.4. A UKPHINS (UK Plant Health Interception Notification System) notification should be made upon confirmation of an interception of *Epirix* spp.. UKPHINS is the IT system for recording findings and non-compliance in order to maintain records and to notify other National Plant Protection Organisations (NPPO) of plant health issues.
- 5.5. If all or part of the consignment has been distributed to other premises prior to diagnosis, trace forward and trace back inspections should take place upon suspicion or confirmation of *Epirix* potato flea beetles. Details of recent past and future consignments from the same grower/supplier should also be obtained.

Official action to be taken following the suspicion of an *Epirix* sp. outbreak

- 5.6. Suspected outbreaks will be assessed on a case by case basis. An Outbreak Triage Group (OTG), chaired by the CPHO or deputy and including specialists from APHA, Defra and other organisations, should subsequently be set up to assess the risk and decide on a suitable response. Where appropriate, the OTG will also decide who will be the control authority, and the control authority will then nominate an Incident Controller. An Incident Management Team (IMT) meeting, chaired by the Incident Commander, will follow the CCG. This will produce an Incident Action Plan to outline the operational plan. See the *Defra Contingency Plan for Plant Health in England* for full details.
- 5.7. The OTG will determine an alert status, which will consider the specific nature of the outbreak. These alert levels, in order of increasing severity, are white, black, amber and red (more details on these levels can be found in table 2 of the *Defra Generic Contingency Plan for Plant Health in England*). Due to the potential for spread, if an infestation of *Epirix* potato flea beetles is suspected in a potato field, an amber alert status should be given by the PHSI. An amber alert status refers to a serious plant pest/disease with potential for relatively slow, but extensive geographical spread leading to host death and/or major economic, food security or environmental impacts.

Restrictions on movement of material, equipment and machinery to and from the place of production

- 5.8. Eggs, larvae, pupae and adults can be transferred in soil associated with non-host material, equipment and machinery, and adults can hitch hike on these (EPPO, 2010). Movement of material, equipment and machinery from infested areas (i.e. the field where the beetle was found) should therefore be restricted. However, if movement is necessary, the material, equipment and machinery should be thoroughly cleaned to remove any soil and life stage of *Epitrix* potato flea beetles.

Preliminary trace forward / trace backward

- 5.9. Based on their biology, *Epitrix* potato flea beetles are most likely to be associated with imports of seed and ware potatoes with soil attached (EPPO, 2010). These potatoes are likely to be from the EU (Portugal and Spain). Third countries (including those where the beetles are present) are prohibited from importing potato tubers, and the potatoes are more likely to be for consumption, as seed potatoes undergo a rigorous selection and testing process prior to import. In comparison to Spain, imports of ware potatoes from Portugal have been small in recent years. Potatoes from Portugal are also generally washed prior to export, minimising the risk of any live stages of *Epitrix* potato flea beetles being present. The risk posed by Spanish ware is therefore of more concern. Several consignments of ware potatoes imported from Spain into the UK in 2015 showed *Epitrix* feeding damage, and in three of these consignments of imported washed potatoes, dead larvae were found to be present in the serpentine tunnels under the skin of the tubers. There were also a number of consignments that had come in unwashed and all of these came from areas thought to be free of *Epitrix* potato flea beetles, and so these areas had no official containment or eradication measures. Although, this risk has now been reduced by national measures introduced in 2015 which require potato tubers to be washed from non-demarcated zones of Portugal and Spain.
- 5.10. If an infested consignment is considered as being the source of the suspect outbreak, investigations regarding the origins of infested consignments will be undertaken to locate other related and therefore potentially infested consignments of tubers or plants moving to and from the site. If applicable the relevant NPPO should be contacted. This process is particularly important for propagation or seed potato stocks.

Confirming a new outbreak

How to survey to determine whether there is an outbreak

- 5.11. Information to be gathered by the PHSI on the suspicion of an infestation of *Epitrix* potato flea beetles, in accordance with ISPM 6; guidelines for surveillance (<https://www.ippc.int/en/publications/615/>)
- The origin of the potatoes and seed lot numbers etc.
 - Details of other premises or destinations where the potatoes have been grown or sent, where the beetles may be present.
 - The layout of the premises and surrounding area, including a map of the fields/cropping/ buildings, at risk growers, and details of neighbouring crops, especially any commercial or non-commercial solanaceous crops in fields, allotments, gardens or glasshouses.
 - Details of the crop variety, growth stage and any other relevant information.
 - Description of the surrounding habitat.
 - Level of infestation, including a description of symptoms (as well as photos if possible). These would include shot-holes on foliage, and serpentine tunnels and holes in tubers.
 - The date and time the sample was taken, how it was identified and by whom.
 - Current treatments/controls in place e.g. chemical treatments.
 - Details on the movement of people, equipment, machinery etc. to and from the infested area.
 - Cultural and working practices.
 - If in a store, details of consignments stored next to the suspect consignment.
- 5.12. Further to information gathering, samples of other beetles and tuber feeding damage should be taken to confirm the extent of the infestation e.g. in surrounding lots or fields. This initial survey will be used to determine if it is an isolated finding or an established outbreak (see 5.24-5.28).
- 5.13. If at a processor/packer, any host plants (including any tubers, which should be held) should be surveyed on the site or in the immediate vicinity in the summer and again in the following year for signs of pest presence (see 5.15 -5.18). If the site is in a potato growing area, the two potato fields closest to the site and a field on the prevailing wind side should also be surveyed (crops in these fields should be held under notice pending inspection). Waste disposal processes and areas should likewise be inspected.
- 5.14. Finance for the surveys will depend on the individual circumstances of the outbreak, and will be subject to discussion, usually between Defra policy and the PHSI.

Sampling

5.15. Adult *Epitrix* beetles are small (1.5 – 2 mm in length), oval, convex and black, and feed on the foliage of host plants. Because they are difficult to see and jump when disturbed, sweep netting should be used to capture the adults. This would ideally be carried out in places where the foliage of host plants exhibits shot-hole like damage, which is indicative of adult presence. There are two methods of sweep netting, depending on the size of the vegetation:

- 1) Young potato plants and herbaceous vegetation surrounding the crop. To use a sweep net, swing it in an approximately 180° arc such that the net rim strikes the top 6 to 8 inches of vegetation. Hold the net slightly less than vertical so the bottom edge strikes the vegetation before the top edge. This will facilitate getting the insects into the net. A common practice is to take a sweep from right to left, walk a step, and take another sweep, left to right (see https://www.youtube.com/watch?v=c5dVt3n1_EE for an example). Do this quickly as *Epitrix* beetles drop off the vegetation when they see you approach. Try to avoid chopping the tops of the plants and filling your net with vegetation. After taking the desired number of sweeps (about four or five), grasp the net bag with a hand at about the mid-point. Shake the bag so that most of the insects fall to a bottom corner of the bag, quickly put an open plastic tube into the corner and pour in the insects caught. Seal the tube and label it. Some insect collectors prefer to put their head into the net and only catch specific target insects using an electronic aspirator or pooter. Take the tube containing the insect catch to the laboratory and cool it in a fridge before opening it and examining the contents. *Epitrix* can jump and move surprisingly fast, particularly in warm conditions, and can therefore be difficult to collect in the field. You may need to practice and modify your technique to suit the local circumstances. Do not sweep wet vegetation.
- 2) Larger potato plants. The sweep net can be used like a beating tray if inspecting mature potato plants. Place the net below the plant and strike it sharply with a stick or give it a quick shake. If there are *Epitrix* beetles present, they should fall into the net. Capture them into a plastic tube using the same procedure as described above.

5.16. Larvae normally feed externally and/or just underneath the surface of potato tubers. Tubers should be inspected for characteristic holes, serpentine tunnels, and black splinter lesion feeding damage (deeper holes with a trail of corky material inside the tuber).

5.17. Tuber inspections are best undertaken on tubers during harvesting, grading or in store because a large number of tubers can be inspected more easily than in the growing crop. Tuber symptoms are also likely to be more evident later in the growing

season. Tubers discarded on grading or packing lines should be targeted for inspection.

- 5.18. Following the identification/capture of an adult, pupae, larva, and/or larval feeding damage, the samples should be sent for diagnosis as in point 5.1. Each sample should be labelled with full details of sample number, location (including grid reference if possible) and variety.

Diagnostic procedures

- 5.19. Morphological diagnosis of *Epitrix* potato flea beetles can only be done for adults and requires microscopic examination of genitalia. EPPO diagnostic protocol PM 7/109 (EPPO, 2011a) is recommended for the identification of *E. cucumeris* and *E. tuberosa*. *Epitrix similis* is also included in the protocol, but this species was included prior to the description of *E. papa*. The use of Orlova-Bienkowskaja (2015) for the diagnosis of *E. similis* and *E. papa* is therefore preferred.

Criteria for determining an outbreak

- 5.20. If *Epitrix* potato flea beetles are detected at a location and are not confined to a particular consignment(s) then an outbreak should be declared. For example, if they are identified in a potato field, then this would be classified as an outbreak. However, if they are restricted to recently imported potatoes within a cold store, then this would be classified as an interception. If only symptoms of *Epitrix* potato flea beetles are found, then the outbreak should be treated as suspected until live stages are found.

Official Action to be taken following the confirmation of an outbreak

- 5.21. The scale of the outbreak will determine the size and nature of the management team and action.

Communication

- 5.22. The Incident Management Team will assess the risks and communicate details to the IPPC and EPPO, in accordance with ISPM 17: pest reporting (<https://www.ippc.int/en/publications/606/>), as well as within Government to Ministers, senior officials and other government departments, devolved administrations and agencies (e.g., the Environment Agency) on a regular basis as appropriate; and to stakeholders.

5.23. The Defra pest factsheet to raise awareness of *Epitrix* potato beetles and their symptoms should be distributed to relevant stakeholders in the locality of where the beetles have been found. The pest factsheet can be found on the Plant Health Portal - <https://planthealthportal.defra.gov.uk/plant-health-api/api/pests/3686/notices/6707/documents/4218/document>

Surveillance and demarcated zones

5.24. After an outbreak has been detected, a regulated area should be established that includes:

- An infested zone (i.e. the infested field or crops).
- A buffer zone, which should extend out at least 1 km radius from the edge of the infested zone.

5.25. Areas where machinery used was the same as that for the infested zone, and any other areas where there is a perceived risk should also be surveyed. If resources are limited, priority should be given to areas where there has been movement of large quantities of soil from the infested zone, though this should be agreed by the IMT on a case by case basis.

5.26. All host plants in the infested zone should be visually inspected, samples taken and suspect samples sent for diagnosis (see 5.15-5.18).

5.27. Potato fields neighbouring the infested zone should also be surveyed, and after this has been completed surveys should be extended to potato fields and hosts within the buffer zone.

5.28. The regulated area should be adjusted in response to further findings. If *Epitrix* potato flea beetles are found within a field outside the infested zone, this should subsequently be designated as infested. If beetles are found within uncropped areas outside the infested zone, then any fields directly adjacent to these areas should normally be designated as infested.

Decontamination procedures

5.29. All non-disposable material, equipment and machinery, should be thoroughly cleaned (e.g. using water at high pressure) to remove the pest and any soil. Ideally, the cleaning should be carried out within the demarcated area.

5.30. Any waste (plant or other potentially infested material) should be removed and destroyed (via deep burial, incineration or other appropriate methods prescribed in 5.48-5.50).

Tracing forwards / backwards

5.31. Once other sites that are potentially infested by *Epitrix* potato flea beetles have been identified, these should be inspected as per 5.15-5.18.

Pest Management procedures (based on EPPO national regulatory control system)

Scenario 1: Outbreak in a potato crop at an early stage of growth (i.e. prior to tuber formation) – infested zone

- 5.32. The whole crop should be treated as soon as possible with a foliar insecticide. Foliar insecticides will kill adults above ground, but will have no effect on the eggs, larvae, pupae or adults within the soil. The Defra Risk and Horizon Scanning team should be consulted about the most appropriate treatments.
- 5.33. Sweep nets should be used to sample for adults before and after spraying to assess the efficacy of insecticide treatments (see 5.15). Sweep netting should not be carried out while the crop is still wet with insecticide, so it is best to leave the crop at least a couple of days after spraying.
- 5.34. When the level of adults is low (no beetles caught), the potato haulm should be destroyed (using methods in 5.47) to eliminate the food supply of *Epitrix* potato flea beetles and thereby reduce their ability to survive and multiply.
- 5.35. If the level of adults is high (beetles caught), the potato haulm should **not** be destroyed due to the high risk of adults dispersing to hosts in other fields/sites.
- 5.36. Even if the level of adults is low when the haulm is destroyed, there is still a risk of some adults (including newly emerging adults) spreading further afield. Several rows of crops should therefore be left at the edge of the field to act as trap crops. These should be treated regularly with insecticide and destroyed later in the season.
- 5.37. Any tubers which may have developed early should not be harvested due to the risk of pest dispersal with the tubers and any associated soil. Destruction of tubers in a way that does not risk further spread (e.g. exposure to frost) should be considered, and will be decided by the Incident Management Team.
- 5.38. Host plants, such as weeds, and other plant material in the infested zone, and uncropped areas, field boundaries and hedgerows in the immediate vicinity of the infested zone should be destroyed by herbicide or mechanical means.

Scenario 2: Outbreak in a growing potato crop (tubers have developed) – infested zone

- 5.39. This is the same as for scenario 1, but because tubers are present for the *Epitrix* potato flea beetle larvae to feed on, the benefit of destroying the potato haulm to remove the food supply for larvae is less clear cut. Furthermore, early destruction of the plant haulm may cause the dispersal of adults, which may be present in higher numbers at this stage of the season. The decision therefore will need to be made on a case by case basis.
- 5.40. The crop should not be harvested because of the risk of spread. Measures should be taken to reduce potential volunteer problems in the following year. This may include the application of a herbicide to the growing crop to prevent sprout development of progeny tubers. Any volunteer potato plants that grow in the field should be destroyed soon after emergence.

Scenario 3: Measures to be taken in the case of detection of infestation in tubers after harvest (e.g. in a grower's potato store or during processing/packaging when the tubers can be traced back to a known crop or farm)

- 5.41. The following should be designated as infested:
- The lot from which the sample was taken and lots that the *Epitrix* potato flea beetles may have spread to.
 - The waste from the infested lot, such as soil and processed waste.
 - The equipment and other articles (e.g. machinery and packing material) which have been in contact with the lot.
 - The field where the lot was grown.
- 5.42. As in point 5.24, a buffer zone should be created that extends out to 1km from the infested field and store.
- 5.43. Areas where machinery used was the same as that for the infested zone, and any other areas where there is a perceived risk should also be surveyed. If resources are limited, priority should be given to areas where there has been movement of large quantities of soil from the infested zone.
- 5.44. There is the risk that other potato stocks may have become infested after harvest. This could have been through the transfer of soil in grading lines or storage boxes. Any tubers in storage should therefore be inspected for symptoms and for the presence of *Epitrix* potato flea beetles.

Crops growing within the buffer zone (1km around the infested zone) in the year of the outbreak

- 5.45. If no infestation is found in the potato crops growing in the buffer zone following surveillance (5.24-5.28) then they should be treated with a programme of foliar insecticides until harvest and monitored for any sign of *Epitrix* potato flea beetles. Potato tubers should also be inspected during and/or immediately after harvesting.
- 5.46. All potato tubers that are harvested should be washed to remove soil and the pest prior to movement outside of the buffer zone. The potatoes will also only be marketed as ware potatoes and will not be used as farm saved seed.

Disposal plan

Infested growing host crops

- 5.47. The primary means of destruction of potato plants in a field is through herbicide application. Actives that can be used include glyphosate and carfentrazone-ethyl (e.g. Spotlight plus), but the Defra Risk and Horizon Scanning team should be consulted for appropriate treatments.
- Any applications should be made following the advice on the product label and be in accordance with HSE guidance. In some cases there may be a requirement to carry out a Local Environment Risk Assessment for Pesticides (LERAP) depending on the product used and the situation of the finding.
 - Prior to any herbicides being used, the risk posed by the herbicide to people and the potential environment impacts will be assessed. The treatment should only be made if it is determined that there will be no unacceptable risks to people and the environment.

Infested harvested tubers

- 5.48. Adults, pupae and possibly larvae may be present with the harvested tubers, associated soil and plant debris. It is important that all of this material is disposed of safely so as to eliminate the pest. When deciding on the most appropriate method(s) of disposal, factors such as the likelihood of adults being present, the level of handling and transportation required, all need to be taken into account. For all methods, measures need to be taken to ensure that there is no risk of spread during transport and treatment or disposal. Tubers that can be moved safely should be destroyed by incineration at a licensed facility (if in small quantities) or deep burial. Disposal and/or destruction should be under the approval and supervision of the PHSI. If the material has to be moved off the premises, it should be contained within at least one sealed layer, and two layers if possible. Deep burial may be done at an approved landfill site, or on the site or nearby farm, but only in agreement with the local Environment Agency. Incineration must comply with appropriate waste

management regulations, Environment Agency in England, Scottish Environment Protection Agency and Natural Resources Wales.

- 5.49. Other possible methods of destruction for potato tubers should be considered on a case by case basis and include heat sterilization, industrial processing (under official supervision), fermentation and composting, steaming and feeding to animals, and anaerobic digestion (minimum temperature of 55°C for 24 h without interruption with a hydraulic dwell time in the reactor of at least 1 day).
- 5.50. All objects designated as 'infested', such as equipment, machinery, storage facilities that may be contaminated with infested soil or potato material should be thoroughly cleaned (e.g. using water at high pressure) to remove the pest and any soil.

Measures in subsequent seasons

Infested zone

- 5.51. No host or root crop should be planted for at least 2 years and until no volunteer plants have been found for 2 consecutive years. Trap crops planted in the year following the outbreak are exempt from this rule.
- 5.52. Any volunteer plants should be destroyed early in the season following the outbreak. If the population of volunteer plants is low, they can be removed by hand, but if the population of volunteer plants is high, they are best controlled by an application of an effective herbicide (see 5.47). Solanaceous weeds should also be controlled in a similar way.
- 5.53. 'Infested' fields may be maintained in permanent pasture with frequent close cutting or intensive grazing. This option has the advantage of providing effective control of potato volunteers and solanaceous weeds.
- 5.54. Following 2 years without volunteer plants, only ware potatoes should be produced (if potatoes are grown), with the growing crop and harvested tubers inspected for *Epitrix* potato flea beetles. If there are no finds of the pest following this, then seed potatoes can also be produced on the field.

Buffer zone (1km around the infested zone)

- 5.55. Host crops should not be planted outdoors for at least 2 years. Volunteer plants and weeds should be destroyed. Following this period, only ware potatoes should be produced with the growing crop and harvested tubers inspected for *Epitrix* potato flea beetles. If there are no finds of the pest, then either seed or ware potatoes can be produced on the field.

5.56. Host crops planted indoors should be monitored, and if *Epitrix* potato flea beetles found, the site (which may include outdoor fields) should be designated as infested and an appropriate programme of insecticides applied.

6. Criteria for declaring eradication / change of policy

6.1. *Epitrix* potato flea beetles can be declared eradicated (by the Chief Plant Health Officer) in potato after at least three years during which time no *Epitrix* potato flea beetles have been found. These three years must include 2 consecutive years without volunteer plants and then either a year with a ware potato crop or trap potato crop.

7. Evaluation and review of the contingency plan

7.1. This pest specific contingency plan should be reviewed regularly to consider changes in legislation, control procedures, pesticides, sampling and diagnosis methods, and any other relevant amendments.

7.2. Lessons should be identified during and after any *Epitrix* or non-*Epitrix* outbreak, including what went well and what did not. These should be included in any review of the contingency plan leading to continuous improvement of the plan and response to outbreaks.

8. Appendix A

Data sheet for *Epitrix similaris*, *Epitrix tuberis*, *Epitrix cucumeris* and *Epitrix papa*

Identity

PREFERRED SCIENTIFIC NAME	AUTHOR (taxonomic authority)
<i>Epitrix similaris</i>	(Gentner, 1944)
<i>Epitrix tuberis</i>	(Gentner, 1944)
<i>Epitrix cucumeris</i>	(Harris, 1851)
<i>Epitrix papa</i>	(Orlova-Bienkowskaja, 2015)

CLASS: Insecta

ORDER: Coleoptera

SUBORDER: Polyphaga

SUPERFAMILY: Chrysomeloidea

FAMILY: Chrysomelidae

SUBFAMILY: Galerucinae

INTERNATIONALLY USED COMMON NAME/S & INTERNATIONAL LANGUAGE

***Epitrix similaris*:**

Amerikanische Kartoffeldlerdlöhe (German)

Potato flea beetle (English)

Pulguilla de la patata (Spanish)

Altise américaine de la pomme de terre (French)

***Epitrix tuberis*:**

Tuber flea beetle (English)

Altise des tubercules (French)

***Epitrix cucumeris*:**

Amerikanischer Kartoffelerdfloh (German)

Nordamerikanischer Kartoffelerdfloh (German)

Potato flea beetle (English)

Pulguilla del pepino (Spanish)

Pulguilla negra (Spanish)

Altise de la pomme de terre (French)

OTHER COMMON NAMES & COUNTRY WHERE USED

Notes on taxonomy and nomenclature

Epitrix is a large genus, which is composed of nearly 180 species (Doeberl, 2000). *Epitrix similis*, *E. tuberos*, *E. cucumeris* and *E. papa* are part of a group of six beetles that also includes *Epitrix subcrinita* and *Epitrix hirtipennis* (tobacco flea beetle) that are all reported to feed on potatoes.

Biology and ecology

Life history

The basic life cycle of the four *Epitrix* species is similar (see fig. 2) and is described by Eyre and Giltrap (2013). Most is known about the life cycle of *E. tuberos*, with fewer studies on the other three species.




Figure 2. Life cycle of *Epitrix* potato flea beetles.

Adult *Epitrix* beetles overwinter in the soil, nearby to their host plants, and this often occurs in field margins, especially if solanaceous weed hosts are present (1 in fig. 2). Their survival in the soil depends partly on the depth at which they overwinter; *E. tuberis* adults survive well at 20-30 cm (Davis and Landis, 1947), and can survive down to depths of 60 cm (Campbell *et al.*, 1989). Following emergence in spring (or summer depending on location), adults feed on the upper leaf surfaces of the host plants (2). *Epitrix tuberis* has also been observed to move to the lower surface under windy conditions and at night (Bérubé, 2000). Adults usually colonise plants on the outer edges of the potato crop first. Movement onto these plants, and those further inwards, can take place via walking, jumping or flying.

Mating occurs within 24 h of emergence in *E. tuberis*. After 5-6 days, adults lay up to 200 eggs over the next 12-55 days, either singly or in small batches of 11-15 in the soil or at the base of the host plant (3, Campbell *et al.*, 1989; Neilson and Finlayson, 1953). The number of eggs laid by any one adult varies depending on the host plant (Boavida *et al.*, 2013).

After an incubation period of 3-14 days, larvae of *E. tuberis* and *E. cucumeris* hatch from the eggs and feed on the roots and tubers of the host plant for 2-4 weeks (4). Pupation subsequently takes place in the soil, lasting around 4-10 days (5). The adults then emerge and begin feeding on the leaves (6). These adults mate and the life cycle begins once more. Up to 3 generations can occur within a single season. This means that thousands of progeny can potentially be derived from each female. Extensive tuber damage can therefore occur in potato crops where very low populations of adults were present at the start of the season.

Hosts/crops affected

The most significant host of all four *Epitrix* species is potato (*Solanum tuberosum*). A recent host preference study undertaken in Portugal also indicates that a further four plants are hosts for *E. papa* and *E. cucumeris* (Boavida *et al.*, 2013). Both species were demonstrated to feed and reproduce on aubergine (*Solanum melongena*), tomato (*Solanum lycopersicum*), and the weeds, black nightshade (*S. nigrum*) and thornapple (*Datura stramonium*). The highest number of progeny produced by *E. papa* was on *S. nigrum* and potato. *Epitrix cucumeris* likewise showed a greater number of progeny on *S. nigrum* than on the other host plants tested. These *Epitrix* species, along with *E. tuberis*, also regularly feed on other members of the nightshade family (Solanaceae), and occasionally feed on other plant families, and may feed on more species than are currently listed (e.g. sugar beet). It should be noted, however, that feeding does not necessarily imply completion of the life cycle on the species. The full list of plants on which feeding by the four species has been recorded are shown in table 1.

Table 1. Host plants of *Epitrix tuberis*, *E. cucumeris*, *E. similaris* and *E. papa* (marked with an X symbol) (CABI, 2014a, b, c; EPPO, 2010). The hosts on which the species have been confirmed to complete their life cycle are marked with the letter H.

Host plant	<i>E. tuberis</i>	<i>E. cucumeris</i>	<i>E. similaris</i>	<i>E. papa</i>
<i>Alcea rosea</i> (Hollyhock)	X			
<i>Allium cepa</i> (onion)		X		
<i>Armoracia rusticana</i> (horseradish)	X			
<i>Beta vulgaris</i> (beetroot)	X	X		
<i>Brassica oleracea</i> (cabbages, cauliflowers)	X	X		
<i>Capsicum annum</i> (bell pepper)		X		
<i>Capsicum frutescens</i> (chilli)	X			
<i>Chenopodium album</i> (fat hen)	X			
<i>Chenopodium spp.</i> (goosefoots)		X		
<i>Cirsium arvense</i> (creeping thistle)	X			
<i>Cucumis sativus</i> (cucumber)	X	X		
<i>Curcubita pepo</i> (ornamental gourd)		X		
<i>Datura stramonium</i> (thorn apple)		XH		XH
<i>Descurainia pinnata</i> (pinnate tansymustard, USA)	X			
<i>Helianthus annuus</i> (sunflower)		X		
<i>Lactuca sativae</i> (lettuce)	X	X		

<i>Medicago sativa</i> (lucerne)	X			
<i>Nicotiana tabacum</i> (tobacco)	X	X		
<i>Phaseolus</i> (beans)		X		
<i>Phaseolus vulgaris</i> (common bean)	X			
<i>Physalis</i> (groundcherry)	X			
<i>Raphanus sativus</i> (radish)	X			
<i>Ribes rubrum</i> (red currant)	X			
<i>Sinapis arvensis</i> (wild mustard)	X			
<i>Solanum jasminoides</i> (potato vine)				
<i>Solanum lycopersicum</i> (tomato)	X	XH		XH
<i>Solanum melongena</i> (aubergine)	X	XH		XH
<i>Solanum nigrum</i> (black nightshade)	X	XH		XH
<i>Solanum rostratum</i> (prickly nightshade)	X			
<i>Solanum triflorum</i> (cutleaf nightshade, USA)	X			
<i>Solanum trifolium</i> (weed)				
<i>Solanum tuberosum</i> (potato)	XH	XH	XH	XH
<i>Spinacia oleracea</i> (spinach)	X			
<i>Stellaria media</i> (common chickweed)	X			
<i>Taraxacum officinale</i> complex (dandelion)				
<i>Zea mays</i> (maize)		X		

Plant stage affected

The *Epitrix* potato flea beetles affect plants during growth and during tuber production.

Plant parts affected

Leaves, roots and tubers.

Symptoms/signs - description

The adult *Epitrix* potato flea beetles feed on foliage, producing numerous “shot-like” holes that are 1-1.5 mm in diameter.

Larvae burrow beneath the surface of tubers, leading to three main types of damage; holes, serpentine tracks and black splinter lesions (Boavida *et al.*, 2013; see Fig. 1 for serpentine tunnels).

Feeding damage by larvae increases the likelihood of infection from pathogenic bacteria and the development of soft rots. The larvae may also vector *Verticillium dahliae*, *Fusarium coeruleum* and *Thanatephorus cucumeris* (EPPO, 2005).

Morphology

Eggs

Elliptical, off-white and approximately 0.5 mm long x 0.2 mm wide (Fig. 2 stage 3).

Larvae

Worm-like, off-white body, with a brown head capsule. When mature, they measure about 5 mm in length (Fig. 2 stage 4).

Pupae

About 2.5 mm in length and off-white in colour (Fig. 2 stage 5)

Adult

Small (1.5 – 2.0 mm in length), oval, convex and brown or black. Species discrimination requires microscopic examination of genitalia (Fig. 5 and 6; also see EPPO, 2011 for details).

Similarities to other species/diseases/plant damages

Epitrix species are difficult to distinguish and identification often requires detailed microscopic examination (of genitalia). This has meant that identification to species level has not been regularly undertaken. As a consequence, reports of damage seen in the field and ascribed to particular species have to be viewed with some caution. Data on the distribution of individual species are therefore also not fully reliable. For example, *Epitrix*

papa, a newly described species, was previously misidentified as *E. similaris* in Portugal and Spain (Orlova-Bienkowskaja, 2015). Molecular diagnostic tools may improve the situation in the coming years (Germain *et al.*, 2013).



Figure 3. Potato flea beetle adults. © EPPO



Figure 4. Potato leaves showing characteristic adult 'shot-hole' feeding damage. © Catherine Chatot, Germicopa SAS, France



Figure 5. *Epitrix cucumeris* adult. © Jean-François Germain. LNPV Montpellier



Figure 6. *Epitrix papa* adult. © Jean-François Germain. LNPV Montpellier



Figure 7. *Epitrix tuberis* larvae. © Agriculture Canada, Ottawa, Bugwood.org

Detection and inspection methods

Leaves can be visually inspected for adult beetles and characteristic “shot-like” holes, whilst potato tubers can be inspected for holes, serpentine tracks and black splinter lesions, which are indicators of *Epitrix* larvae.

History of introduction/spread

In 2004, potato tuber damage indicative of *Epitrix* was observed in north Portugal, near Porto. *Epitrix cucumeris* and a second *Epitrix* species were confirmed across Portugal in 2008. The second species was first identified as *E. similaris*, but this has now been corrected, and was confirmed as *E. papa* in 2015. This new species has now been reported from the Portuguese counties of Alentejo (2017), Calheta (2018), Machico (2018), Ponta Do Sol (2018), Porto Moniz (2018) and São Vicente (2018). *Epitrix papa* has also been recorded in Spain, in Galicia (2010 – date of first record), Asturias (2014), and Andalucia (2015).

Interestingly, *Epitrix cucumeris* is also found in the Azores, where it has been since at least 1979, and Madeira, but no damage on potato tubers has been reported.

E. similaris and *E. tuberis* are reported as absent from Europe. All species are absent from the UK.



Figure 12. Locations and regions in which *E. cucumeris* and *E. papa* have been detected in Portugal and Spain.

Distribution

Epitrix similaris




Figure 8. *Epitrix similaris* distribution as of July 2021. (Source: EPPO Global database). The link below provides up to date distribution data.

<https://gd.eppo.int/taxon/EPIXSI/distribution>

Epitrix tuberos



Figure 9. *Epitrix tuberos* distribution as of July 2021. (Source: EPPO Global database). The link below provides up to date distribution data.

<https://gd.eppo.int/taxon/EPIXTU/distribution>

Epitrix cucumeris




Figure 10. *Epitrix cucumeris* distribution as of July 2021. (Source: EPPO Global database). The link below provides up to date distribution data.

<https://gd.eppo.int/taxon/EPIXCU/distribution>

Epitrix papa




Figure 11. *Epitrix papa* distribution as of July 2021. (Source: EPPO Global database). The link below provides up to date distribution data.

<https://gd.eppo.int/taxon/EPIXPP/distribution>

Phytosanitary status

Epitrix similaris was an A2 EPPO listed pest, but was removed from the list in 2016, following the recognition the description of *E. papa*.

Epitrix tuberis is a GB quarantine pest, on the EPPO A1 list, the East Africa A1 list, the Southern Africa A1 list, the Jordan A1 list, the Kazakhstan A1 list, the Georgia A1 list, the Russia A1 list, the Turkey A1 list, the Ukraine A1 list, and is a quarantine pest in Morocco, Tunisia, Mexico and Israel.

Epitrix cucumeris is a GB quarantine pest, on the EPPO A2 list, the Egypt A1 list, the Bahrain A1 list, the Jordan A1 list, the Kazakhstan A1 list, the Turkey A1 list, the Ukraine A1 list, and is a quarantine pest in Morocco, Tunisia and Jordan.

Epitrix papa is a GB quarantine pest, on the EPPO A2 list and the Ukraine A1 list.

Means of movement and dispersal into the UK

Although adult beetles can fly, there is little evidence confirming long distance spread of these species through natural dispersal. Instead, the main pathway is likely to be via human transport, principally the movement of seed and ware potatoes with soil attached. Other, less likely, pathways include the movement of free soil, the movement of soil attached to other host and non-host plants, and the movement of soil attached to machinery, from infested areas.

Control

Chemical control

Once established, *Epitrix* is primarily controlled using a programme of foliar applied insecticide treatments. These control adults but have no significant impact on the larvae or pupae in the soil. Although granular insecticides applied to the soil can also be used as a means of targeting larvae and pupae, these have so far provided inadequate protection and are expensive (Finlayson *et al.*, 1979). They are also not currently approved in the UK.

Foliar spray timing is critical, especially early on in the season when the first generation adults emerge. Early spraying ensures the adults are suppressed before they lay their eggs, reducing the number of second generation adults that emerge later in the season.

Of the insecticides used, synthetic pyrethroids are particularly effective as foliar sprays (Vernon and Mackenzie, 1991). An early season application of one of these insecticides can reduce tuber damage by > 70%. Later in the season, other insecticides that are currently sprayed against other insect pests, such as the Colorado beetle, provide control.

Although they have the potential to overwinter in the centre of potato fields (Vernon and Thomson, 1991), *Epitrix* potato flea beetles tend to concentrate at the field perimeter. *Epitrix* potato flea beetles can therefore be controlled sufficiently by spraying the outer rows of the potato crop, as is the case for *E. tuberis* in Canada.

Reliance on insecticides means that control in organic crops will be very difficult. The development of resistance is also a concern. *Epitrix tuberis* has previously shown resistance to DDT (Kring, 1958) and organochlorines (Campbell and Finlayson, 1976). Additionally, there are far fewer insecticides available in the EU than there are in North America.

Field monitoring/economic threshold levels

In the USA and Canada, treatment thresholds have been developed to help decision making on when the early treatments should be applied. These are based on visual inspections of potato plants in the outer rows of the crop for adult beetles, usually 10 per row, over a period of 5-20 seconds per plant. If plants exceed 30 cm, sampling is usually done by sweep netting, in which 10 adjacent circular sweeps are carried out to give a sample of 100 plants. Economic thresholds vary depending on the *Epitrix* species, location and plant variety. For example, the economic threshold for *E. cucumeris* in Manitoba is 65 beetles per plant for the variety Norland and 190 per plant for Russet Burbank (Senanayake *et al.*, 1993). No thresholds have been developed for treatment in Portugal or Spain. Instead, growers inspect crops and usually apply sprays when adults and/or the first signs of foliar damage are found in potato crops.

Cultural controls and sanitary methods

Growing non-host crops (rotation) can help reduce the build-up of *Epitrix* beetle populations; Kabaluk and Vernon (2000) found that the cost of controlling *E. tuberis* was over £12 greater per hectare on fields that had potatoes planted for the last 3 years, as compared with a field that did not have potatoes in the previous year. Delaying the planting of potatoes until later in the season might also be effective. This forces *Epitrix* beetles to first find alternative hosts that are likely poorer in quality. Further controls include the removal of potential hosts (e.g. solanaceous weeds) over winter and during the season, and the careful disposal of reject potatoes.

Impacts

Economic impact

In 1904, when *E. tuberis* was first recorded in North America, losses amounted to \$250,000 (Gentner, 1944). Yield losses of 22-25% and 43% have subsequently been reported on Prince Edward Island and Canada, respectively (Stewart and Thomson, 1989). The level of these losses represents a considerable danger to the EU, which produces 62 million tons of potatoes each year, with sales exceeding £2.5 billion per

annum (Schwartzmann, 2010). In Portugal, the greatest losses occurred in the first few years before *Epitrix* potato flea beetles had been identified as the cause of the problem. Consignments of tubers showing damage were rejected by potato packers. In recent years, damage has been reduced in Portugal through the application of one or two insecticide sprays early in the season, in addition to those applied later for Colorado beetle control (Oliveira pers. comm. 2010). These treatments do not eliminate tuber damage, but reduce it to a level acceptable for the quality requirements of the country's internal market, which tend to be less stringent than those required in the UK and northern Europe.

Control of *Epitrix* potato flea beetles with insecticides requires precise timing, and it is likely that this will not always be met. Economic losses, beside those relating to chemical treatment, will therefore still occur. In the UK, *Epitrix* potato flea beetles could affect the marketability of ware and seed potatoes, particularly the latter, which represent a high risk pathway. Potato growers may therefore be very reluctant to buy seed if there is any perceived risk associated with it. Estimated losses following the establishment of *Epitrix* potato flea beetles could be as high as £10-41 million per year, despite the use of insecticides.

Environmental impact

Epitrix potato flea beetles primarily feed on plants within the Solanaceae family, and occasionally on plants from other families, such as cabbages and cucumbers. Although wild solanaceous species (and possibly other plant groups) may be targeted and may act as reservoirs for the beetle, it is unlikely that the environmental impact will be severe. The impact is therefore largely economic and derived from the direct effects of the beetles on crop yield and quality of potato. However, where *Epitrix* potato flea beetles are present, insecticides are used to control them, and this may have a potentially significant effect on non-target arthropods.

9. Appendix B

Statutory Action

- For findings highlighted in Green, a UKPHINS notification will be made if live beetles/larvae are detected.
- For findings in yellow, no UKPHINS notification will be made, except in cases where dead larvae are found, but an informal notification will be sent bilaterally to Spain or Portugal, as appropriate.

Table 6. Statutory action taken against potato tubers imported from Portugal and Spain.

Origin	Action – live larvae or beetles	Action – feeding damage/dead larvae on washed potatoes <0.1% soil	Action – feeding damage/dead larvae on brushed potatoes <0.1% soil	Action - feeding damage/dead larvae on unwashed/unbrushed potatoes (or washed/brushed with >0.1% soil)	Action – washed potatoes with <0.1% soil	Action – unwashed/brushed potatoes with >0.1% soil	Action – brushed potatoes with <0.1% soil	Action - non-notified consignments	Action - Incorrect or Absent PP
Spain mainland – other than Demarcated Areas	Destruction or Re-export	No action unless in exceptional circumstances (i.e. suspicion of pest presence) but may lead to increased	Destruction or Re-export	Destruction or Re-export	No Action	Destruction or Re-export	Destruction or Re-export	Destruction or Re-export	N/A

Origin	Action – live larvae or beetles	Action – feeding damage/dead larvae on washed potatoes <0.1% soil	Action – feeding damage/dead larvae on brushed potatoes <0.1% soil	Action - feeding damage/dead larvae on unwashed/unbrushed potatoes (or washed/brushed with >0.1% soil)	Action – washed potatoes with <0.1% soil	Action – unwashed/brushed potatoes with >0.1% soil	Action – brushed potatoes with <0.1% soil	Action - non-notified consignments	Action - Incorrect or Absent PP
		inspection rates							
Portugal and Spain – Demarcated Areas	Destruction or Re-export	No action unless in exceptional circumstances (i.e. suspicion of pest presence) but may lead to increased inspection rates	No action unless in exceptional circumstances (i.e. suspicion of pest presence) but may lead to increased inspection rates	Destruction or Re-export	No Action	Destruction or Re-export	No Action	Destruction or Re-export	Destruction or Re-export

Origin	Action – live larvae or beetles	Action – feeding damage/dead larvae on washed potatoes <0.1% soil	Action – feeding damage/dead larvae on brushed potatoes <0.1% soil	Action - feeding damage/dead larvae on unwashed/unbrushed potatoes (or washed/brushed with >0.1% soil)	Action – washed potatoes with <0.1% soil	Action – unwashed/brushed potatoes with >0.1% soil	Action – brushed potatoes with <0.1% soil	Action - non-notified consignments	Action - Incorrect or Absent PP
Portugal – other than Demarcated Areas, Spain – Balearic Islands, and other EU countries where the beetles are not present	Destruction or Re-export	No action unless in exceptional circumstances (i.e. suspicion of pest presence) but may lead to increased inspection rates	Destruction or Re-export	Destruction or Re-export	No Action	No Action	No Action	Destruction or Re-export	N/A

10. References

- Antonelli, A. L. and Davidson, R. M. jr** (1991) Potato flea beetles: biology and control. Extension Bulletin – Cooperative Extension, College of Agriculture & Home Economics, Washington State University [Online] No. EB1198:2. Available: <http://whatcom.wsu.edu/ag/documents/seedpotatoes/eb1198e.pdf>. [Accessed: 13/11/2014].
- Arnett, R. H.** (1985) American insects: A handbook of the insects of America north of Mexico. New York, USA: Van Nostrand Reinhold.
- Bérubé, C.** (2000) Potato IPM: tuber flea beetle [online]. Available: <http://nanaimo.ark.com/~cberube/tfb.htm>.
- Boavida, C., Giltrap, N., Cuthbertson, A. G. S. and Northing, P.** (2013) *Epitrix similaris* and *Epitrix cucumeris* in Portugal: damage patterns in potato and suitability of potential hosts plants for reproduction. *EPPO Bulletin*. **43**, 323-333.
- Bousquet, Y.** (1991) Checklist of beetles of Canada and Alaska, Ottawa, Canada: Research Branch, Agriculture Canada Publication.
- CABI** (2014a). *Epitrix cucumeris* In: Crop Protection Compendium. Wallingford, UK: CAB International. www.cabi.org/cpc.
- CABI** (2014b). *Epitrix similaris* In: Crop Protection Compendium. Wallingford, UK: CAB International. www.cabi.org/cpc.
- CABI** (2014c). *Epitrix tuberis* In: Crop Protection Compendium. Wallingford, UK: CAB International. www.cabi.org/cpc.
- Campbell, C. J. and Finlayson, D. G.** (1976) Comparative efficacy of insecticides against tuber flea beetle and aphids in potatoes in British Columbia. *Canadian Journal of Plant Science*. **56**, 869-875.
- Campbell, J. M., Sarazin, M. J. and Lyons, D. B.** (1989) Canadian beetles (Coleoptera) injurious to crops, ornamentals, stored products, and buildings. Ottawa, Ontario, Canada: Research Branch, Agriculture Canada.
- Davis, E. W. and Landis, B. J.** (1947) Overwintering of potato flea beetles in the Yakima Valley. *Journal of Economic Entomology*. **40**, 821-824.
- Doeberl, M.** (2000) Contribution to the knowledge of the genus *Epitrix* Foudras, 1860 in the Palearctic region (Coleoptera: Chrysomelidae: Alticinae). *Mitteilungen des Internationaler Entomologischer Verein*. **25**, 1-23.

- EPPO** (2005) Data sheets on quarantine pests *Epitrix cucumeris*. *EPPO Bulletin*. **35**, 363-364.
- EPPO** (2009) Generic elements for contingency plans PM 9/110. *EPPO Bulletin*. **39**, 471-474.
- EPPO** (2010) Pest Risk Analysis for *Epitrix* Species Damaging Potato Tubers [Online]. Available: http://www.eppo.int/QUARANTINE/Pest_Risk_Analysis/PRA_intro.htm [Accessed: 13/11/2014].
- EPPO** (2011a) Diagnostic standard for *Epitrix cucumeris*, *E. similaris*, *E. tuberis* PM 7/109. *EPPO Bulletin*. **39**, 501–508.
- EPPO** (2011b) Archives of the EPPO reporting service [online]. Available: http://archiveseppo.org/EPPOReporting/Reporting_Archives.htm. [Accessed: 10/11/2014].
- EPPO** (2013) PQR database. Paris, France: European and Mediterranean Plant Protection Organization. <http://www.eppo.int/DATABASES/pqr/pqr.htm>. [Accessed 10/11/2014].
- Eyre, D. & Giltrap, N.** (2013). *Epitrix* flea beetles: new threats to potato production in Europe. *Pest Management Science*. **69**, 3-6. Doi:10.1002/ps.3423.
- Finlayson, D. G., Wilkinson, A. T. S. and MacKenzie, J. R.** (1979) Efficacy of insecticides against tuber flea beetles, wireworms and aphids in potatoes. *Journal of the Entomological Society of British Columbia*. **76**, 6-9.
- Gentner, L. G.** (1944) The black flea beetles of the genus *Epitrix* commonly identified as *cucumeris* (Harris) (Coleoptera: Chrysomelidae). *Proceedings of the Entomological Society of Washington*. **46**, 137–149.
- Germain, J-F, Chatot, C., Meusnier, I., Artige, E., Rasplus, J-Y. and Cruaud, A.** (2013) Molecular identification of *Epitrix* potato flea beetles (Coleoptera: Chrysomelidae) in Europe and North America. *Bulletin of Entomological Research*. **103**, 354-362.
- IPPC** (2006) ISPM No. 9 Guidelines for Pest Eradication Programmes. In: International Standards for Phytosanitary Measures. pp. 109-118.
- Kabaluk, J. T. and Vernon, R. S.** (2000) Effect of crop rotation on populations of *Epitrix tuberis* (Coleoptera: Chrysomelidae) in potato. *Journal of Economic Entomology*. **93**, 315-322.
- Kirk, V. M.** (1975) A list of beetles of South Dakota. Brookings, USA: Agricultural Experiment Station, Technical Bulletin No. 42, 116.
- Kring, J. B.** (1958) Feeding behaviour and DDT resistance of *Epitrix cucumeris*. *Journal of Economic Entomology*. **53**, 823-828.

- Neilson, C. L. and Finlayson, D. G.** (1953) Notes on the biology of the tuber flea beetle, *Epitrix tuberis* Gentner (Coleoptera: Chrysomelidae), in the interior of British Columbia. *The Canadian Entomologist*. **85**, 31–32.
- Orlova-Bienkowskaja, M. J.** (2015) *Epitrix papa* sp. n. (Coleoptera: Chrysomelidae: Galerucinae: Alticini), previously misidentified as *Epitrix similaris*, is a threat to potato production in Europe. *European Journal of Entomology*. **112**, 824-830.
- Schwartzmann, M.** (2010) Potato – a world production, a European business [Online]. Proceedings of the Twelfth Euroblight Workshop. PPO-Special Report. 14, 11-16. Available from:
http://130.226.173.223/euroblight/Workshop/2010Arras/Proceedings/Page11_16_Schwartzmann_web.pdf. [Accessed: 13/11/2014].
- Seeno, T. N. and Andrews, F. G.** (1972) Alticinae of California, Part 1: *Epitrix* spp. (Coleoptera: Chrysomelidae). *Coleopterists Bulletin*. **26**, 53-61.
- Senanayake, D. G., Pernal, S. F. and Holliday, N. J.** (1993) Yield responses of potatoes to defoliation by the potato flea beetle (Coleoptera: Chrysomelidae) in Manitoba. *Journal of Economic Entomology*. **85**, 1527–1533.
- Stewart, J. G. and Thomson, L. S.** (1989) The spatial distribution of spring and summer populations of adult potato flea beetles, *Epitrix cucumeris* (Harris) (Coleoptera: Chrysomelidae), on small plots of potatoes. *The Canadian Entomologist*. **121**, 1097–1101.
- Vernon, R. S. and Mackenzie, J. R.** (1991) Evaluation of foliar sprays against the tuber flea beetle, *Epitrix-tuberis* Gentner (Coleoptera, Chrysomelidae), on potato. *Canadian Entomologist*. **123**, 321-331.
- Vernon, R. S. and Thomson, D. R.** (1991) Overwintering of tuber flea beetles, *Epitrix tuberis* Gentner (Coleoptera: Chrysomelidae), in potato fields. *The Canadian Entomologist*. **123**, 239-240.
- Wilcox, J. A.** (1975) Check list of the beetles of Canada, United States, Mexico, Central America and the West Indies. Latham, New York, USA: Biological Research Institute of America.

11. Authors and reviewers

Authors:

Original: Matthew Everatt & Neil Giltrap (Defra) (2017)

Revised by: Simon Honey (Defra) (2022)

Reviewers:

Janet Dixon (Defra)

Dominic Eyre (Defra)

Joe Ostoja-Starzewski (Fera)

Ian Sanders (APHA)