



Department
for Environment
Food & Rural Affairs

Pest specific plant health response plan: Outbreaks of *Leptinotarsa decemlineata* (Colorado beetle) on potato crops



Figure 1. Adult *Leptinotarsa decemlineata* feeding on a potato leaf. Image Courtesy of Fera-Science Limited © Copyright Fera-Science Limited 2016

© Crown copyright 2017

You may re-use this information (not including logos) free of charge in any format or medium, under the terms of the Open Government Licence. To view this licence, visit www.nationalarchives.gov.uk/doc/open-government-licence/ or write to the Information Policy Team, The National Archives, Kew, London TW9 4DU, or e-mail: psi@nationalarchives.gsi.gov.uk

This document/publication is also available on our website at:

<https://planthealthportal.defra.gov.uk/pests-and-diseases/contingency-planning/>

Any enquiries regarding this document/publication should be sent to us at:

The UK Chief Plant Health Officer

Department for Environment, Food and Rural Affairs

Room 11G32

Sand Hutton

York

YO41 1LZ

Email: plantpestrisks@defra.gsi.gov.uk

Contents

1. Introduction and scope	5
2. Anticipate.....	5
3. Assess.....	6
4. Prepare.....	6
5. Response	8
Official action to be taken following the suspicion or confirmation of <i>Leptinotarsa decemlineata</i> on imported plants, including seeds.....	8
Holding consignments at ports of entry, including packhouses	8
Official action to be taken following the suspicion of <i>Leptinotarsa decemlineata</i> outbreak	9
Restrictions on movement of material, equipment and machinery to and from the place of production.....	9
Preliminary trace forward/trace backward	10
Confirming a new outbreak.....	10
How to survey to determine whether there is an outbreak	10
Sampling	11
Diagnostic procedures.....	11
Criteria for determining an outbreak.....	11
Official action to be taken following the confirmation of an outbreak.....	11
Communication.....	12
Surveillance and demarcated zones.....	12
Decontamination procedures.....	12
Tracing forwards/backwards.....	12

Pest management procedures	13
Disposal plan	14
Measures in subsequent seasons.....	15
6. Criteria for declaring eradication / change of policy	16
7. Evaluation and review of the contingency plan.....	16
8. Appendix A	17
Data sheet for <i>Leptinotarsa decemlineata</i>	17
Identity	17
Notes on taxonomy and nomenclature	18
Biology and ecology	18
Symptoms/signs - description.....	20
Morphology.....	20
Similarities to other species/diseases/plant damages	21
Detection and inspection methods.....	21
Distribution	21
History of introduction/spread	24
Phytosanitary status	25
Means of movement and dispersal.....	26
Control.....	26
Impacts.....	28
9. References	29
10. Authors and reviewers	34
Authors:.....	34
Reviewers:	34

1. Introduction and scope

- 1.1. This pest specific response plan has been prepared by the Defra Risk and Horizon Scanning team. It describes how the Plant Health Service for England will respond if an infestation of *Leptinotarsa decemlineata* is discovered on potato (*Solanum tuberosum*).
- 1.2. This document will be used in conjunction with the Defra *Generic Contingency Plan for Plant and Bee Health in England* (https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/593508/generic-contingency-plan-plant-bee-health-england.pdf), which gives details of the teams and organisations involved in pest response in England, and their responsibilities and governance. It also describes how these teams and organisations work together in the event of an outbreak of a plant health pest.
- 1.3. The aim of this response plan is to facilitate the containment and eradication of *L. decemlineata*.

2. Anticipate

- 2.1. *Leptinotarsa decemlineata* was first recorded in 1811 in the USA by Thomas Nuthall and was first recorded damaging potato plants in 1859, 100 miles west of Omaha, Nebraska (Pope and Madge, 1984). The beetle has since spread across the USA, and moved into Canada, Central America, Europe and Asia (EPPO, 2016). It first established in Europe in Bordeaux, France, in 1922, and is now present in most European countries (Fig. 6; EPPO, 2016). However, it has yet to establish in the UK (EPPO, 2016).
- 2.2. Larvae and adults of *L. decemlineata* feed on the foliage of potato plants, and can completely strip plants of their leaves if they are left uncontrolled. In some EPPO countries, the resulting yield losses have been as high as 50% (EPPO, 1997). Large economic losses (millions of dollars per year) have also been recorded in the USA and China (Grafius, 1997; Liu *et al.*, 2012).
- 2.3. Since 1952, there have been two outbreaks of *L. decemlineata* in the UK, one in 1976 and one in 1977. Both outbreaks were eradicated shortly after detection. *Leptinotarsa decemlineata* is also regularly intercepted on leafy vegetables, salad leaves, fresh herbs and grains, as well as on potato (Fera interception records), from continental Europe.
- 2.4. Eggs, larvae and adults of *L. decemlineata* are mainly associated with potato plants. However, solanaceous plants are prohibited from entering the UK from third countries, and potato plants entering from the EU are tested for harmful organisms (see points 4.4 – 4.12).

The entry of *L. decemlineata* on potato plants is therefore unlikely. Larvae, pupae and adults are also associated with seed and ware potatoes and associated soil. Due to the rigorous selection and testing process prior to import on seed potatoes, ware potatoes and any associated soil are more of a risk. These potatoes are likely to be from Portugal and Spain, with the latter accounting for most of the imports. This pathway is mitigated in part by the emergency and national measures used against *Epitrix* potato flea beetles, which ensures that ware potatoes coming from *Epitrix* demarcated areas in Portugal and Spain must be brushed and/or washed, and that ware potatoes coming from non-demarcated areas of Spain must be washed (see points 4.9 – 4.10). It is for this reason that we are most likely to see *L. decemlineata* “hitch-hiking” on unregulated plants and produce (that may be from nearby to infested potato fields, where beetles may overwinter or use for harbourage), and this has been the case over the last few years.

3. Assess

- 3.1. The beetle currently has a mitigated risk register rating of 20, which is reviewed as and when new information becomes available (<https://secure.fera.defra.gov.uk/phiw/riskRegister/viewPestRisks.cfm?cslref=4840>).
- 3.2. A pest risk analysis, commissioned by the Norwegian Agricultural Inspection Service, was carried out by the Plant Protection Centre in Norway (http://www.mattilsynet.no/planter_og_dyrking/planteskadegjorere/leptinotarsa_decemlineata_1996.4002/binary/Leptinotarsa%20decemlineata%20-%201996).

4. Prepare

- 4.1. *Leptinotarsa decemlineata* is an IB EU listed pest. Countries with protected zone status include **Cyprus**, **Finland** (the districts of Åland, Turku, Uusimaa, Kymi, Häme, Pirkanmaa, Satakunta), **Ireland**, **Malta**, **Portugal**, **Spain** (Ibiza and Menorca), **Sweden** (Blekinge, Gotland, Halland, Kalmar, Skåne) and the **UK**.
- 4.2. *Leptinotarsa decemlineata* is an A2 listed pest in the EPPO region and is therefore recommended for regulation by EPPO member countries.
- 4.3. The Plant Health Service should be aware of the measures described in this plan and be trained in responding to an outbreak of *L. decemlineata*. It is important that capabilities in detection, diagnosis, and risk management are available.

Plants of Solanaceae

- 4.4. Plants of Solanaceae intended for planting, other than seeds, ware and seed potatoes, and other stolon- or tuber-forming plants for planting, are prohibited from third countries other than European and Mediterranean countries (Annex III, Council Directive 2000/29/EC).
- 4.5. Prohibited solanaceous plants can be imported and held under a plant health licence in quarantine conditions (usually for research purposes). Once work on the plants has been completed, destruction of the plants is normally required. However, given adequate testing,

the plants can, in some cases, be released from the terms of the licence if they are shown to be free of pests and pathogens.

- 4.6. Plants of stolon- or tuber-forming species of *Solanum* intended for planting (other than *Solanum tuberosum*) must also be tested and found free of harmful organisms (Annex IV part A section 2, Council Directive 2000/29/EC).

Potato

- 4.7. The import of *Solanum tuberosum* tubers for propagation (seed potatoes) into the EU from third countries other than Switzerland is prohibited under Annex III, Council Directive 2000/29/EC. There is a derogation to bring in tubers for trial, scientific or varietal selection purposes (Commission Directive 2008/61/EC), but they must be placed under quarantine conditions and tested for pests before being used for propagation purposes. There is also derogation for Canadian seed potatoes. This allows Greece, Spain, Italy, Cyprus, Malta and Portugal to import, under specified conditions, seed potatoes originating from certain provinces of Canada:

<http://eurlex.europa.eu/legal-content/EN/TXT/HTML/?uri=CELEX:32014D0368&from=EN>

<http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32011D0778&from=EN>

- 4.8. Ware potatoes and their hybrids are prohibited from third countries, with the exception of Algeria, Egypt, Israel, Libya, Morocco, Syria, Switzerland, Tunisia and Turkey, and other European third countries which are recognised as being free from *Clavibacter michiganensis* ssp. *sepedonicus*, or in which provisions recognised as equivalent to those of the EU territory for mitigating *C. michiganensis* spp. *sepedonicus* have been fulfilled (Annex III, Council Directive 2000/29/EC).
- 4.9. Movement of potato tubers within the EU from *Epitrix* demarcated to non-demarcated areas must be accompanied by a plant passport that is issued as per Commission Directive 92/105/EEC. The potato tubers must also meet the following conditions:
- That they have been grown in a registered place of production as per Commission Directive 92/90/EEC, or by a registered producer or moved from a warehouse or dispatching centre registered as per Commission Directive 93/50/EEC.
 - That they have been washed or brushed before export so that no more than 0.1% of soil remains if they have been imported from a demarcated area in Portugal or Spain.
 - That they are packaged in material that is clean.
- 4.10. From non-demarcated areas for *Epitrix*, the potatoes must be washed before export from mainland Spain in order to kill or remove any *Epitrix* which might be present.
- 4.11. Within the EU, potato tubers for planting must come from advanced breeding selections, have been produced within the EU, been maintained under appropriate conditions, and

tested for harmful organisms using appropriate methods (Annex IV part A section 2, Council Directive 2000/29/EC).

- 4.12. Seed potatoes can only be marketed in the EU if they meet the requirements of Council Directive 2002/56/EU (includes information on certification schemes to ensure healthy plant material).
- 4.13. In response to other pests and diseases of potato, such as *Ralstonia solanacearum* (brown rot) and *Epitrix* potato flea beetles, seed and ware potatoes are surveyed. Any evidence of *Leptinotarsa decemlineata* and its symptoms are more likely to be picked up.

5. Response

Official action to be taken following the suspicion or confirmation of *Leptinotarsa decemlineata* on imported plants, including seeds

Holding consignments at ports of entry, including packhouses

- 5.1. If *L. decemlineata* is suspected by the Plant Health and Seeds Inspectorate (PHSI) to be present in a consignment moving in trade, the PHSI must hold the consignment until a diagnosis is made. Ideally, the consignment should be placed in a sealed cold store and any opened containers should be resealed. Samples should be sent to Fera Science Ltd., National Agri-Food Innovation Campus, Sand Hutton, York, YO41 1LZ (01904 462000) in a sealed bag or container, within at least two other layers of containment, which are not liable to be crushed during transit (no food should be included, but a drop of water can be provided).
- 5.2. When an infestation of *L. decemlineata* is confirmed, the PHSI should advise the client of the action that needs to be taken by way of an official plant health notice. The consignment should be destroyed by either incineration or deep burial (as in 5.58), or re-exported in a sealed container. For some consignments, sorting is permitted (see consignment specific standard operating procedures (SOPs)). If it is not possible to organise destruction by deep burial or incineration, suitable alternatives should be discussed with the Defra Risk and Horizon Scanning team.
- 5.3. Where there is a high risk of escape before destruction or re-export, fumigation may be used under guidance from the Defra Risk and Horizon Scanning team. Fumigation is also preferred over destruction or re-export for timber.
- 5.4. For consignments where beetles are restricted to the outside e.g. on wood packaging, action only needs to be taken to destroy the beetles themselves.
- 5.5. When only dead beetles are found, sorting of the consignment may be permitted. It should be noted that live beetles can enter a state of torpor at low temperatures, becoming immobile, and therefore giving the appearance of being dead. Thus, beetles should be kept at room temperature for several hours to confirm their status.

- 5.6. As appropriate, any host plants (as listed in table 1 and including any potato tubers) should be surveyed by the PHSI for signs of pest presence in the immediate vicinity (the premise in which the pest is found) of any outbreak if this occurs during the growing season and again during the following growing season e.g. in mid-summer (see 5.20-5.22). If the site is in a potato growing area, the two potato fields closest to the site and a field on the prevailing wind side should also be visually surveyed during the growing season. This is a guide and may differ depending on the specific characteristics of the area. Waste disposal processes and areas should also be inspected.
- 5.7. A Europhyt notification should be made upon confirmation of an infestation of live *L. decemlineata*.
- 5.8. Where part of the consignment has been distributed to other premises prior to a confirmed diagnosis, trace forward inspections should take place on suspicion of the presence of *L. decemlineata*. Details of future and previous consignments from the same grower/supplier should also be obtained from the supplying country NPPO.
- 5.9. A pest alert to raise awareness of *L. decemlineata* and its symptoms should be distributed to packers/processors and importers where *L. decemlineata* has been found, and to those in the local area and those associated with the infested premises.

Official action to be taken following the suspicion of *Leptinotarsa decemlineata* outbreak

- 5.10. A Contingency Core Group (CCG), chaired by the Chief Plant Health Officer (CPHO) or their deputy and including specialists from APHA, Defra and other organisations, should be set up to assess the risk and decide on a suitable response at strategic and operation levels. This may include gathering more information on the suspected outbreak, such as the completion of the outbreak assessment, notification of ministers and senior officials, and agreeing a communications strategy. The CCG will also decide who will be the control authority, and the control authority will then nominate an incident commander. An Incident Management Team (IMT) meeting, chaired by the Incident Commander, will subsequently convene to produce an Incident Action Plan (IAP) to outline the operational plan. See the Defra *Generic Contingency Plan for Plant and Bee Health in England* for full details.
- 5.11. The CCG will set an alert status, which will take into account of the specific nature of the outbreak. Under most scenarios, an infestation of *L. decemlineata* suspected in a potato field is likely to be given an amber alert status. An amber alert status refers to a serious plant pest/disease with potential for relatively slow, but extensive geographical spread leading to host death and/or major economic, food security or environmental impacts.

Restrictions on movement of material, equipment and machinery to and from the place of production

- 5.12. Larvae, pupae and adults can be transferred in soil associated with non-host material, equipment and machinery, and adults can hitch hike on these. Movement of material, equipment and machinery between infested and non-infested areas should therefore be restricted. However, if movement is necessary, the material, equipment and machinery

should be thoroughly cleaned at the designated outbreak site to remove any soil and life stage of *L. decemlineata*.

Preliminary trace forward/trace backward

- 5.13. Information obtained regarding the origins of infested consignments should be used to locate other related and therefore potentially infested consignments. Information should also be obtained on the destination to which suspect consignments have been sent. This process is particularly important for propagation or seed potato stocks.
- 5.14. In addition to trace forward investigations relating to seed or ware linked to the outbreak crop we would also need to carry out trace forward / back investigations linked to machinery used on the infested field. Modern farms often share equipment e.g. potato harvesters/planters.

Confirming a new outbreak

How to survey to determine whether there is an outbreak

- 5.15. Information to be gathered by the PHSI on the suspicion of an infestation of *L. decemlineata*, in accordance with ISPM 6; guidelines for surveillance (http://www.acfs.go.th/sps/downloads/13717_ISPM_6_E.pdf):
 - The origin of the potatoes and seed lot numbers etc.
 - Details of other premises or destinations where the potatoes have been grown or sent, where the beetle may be present.
 - The layout of the premises and surrounding area (in relation to potential buffer zones), including a map of the fields/cropping/ buildings, at risk growers, and details of neighbouring crops, especially any commercial or non-commercial solanaceous crops in fields, allotments, gardens or glasshouses.
 - Details of the crop variety, growth stage and any other relevant information.
 - Description of the surrounding habitat.
 - Area and level of infestation, including life stages and a description of symptoms (could take photos). Symptoms would include defoliation and black, sticky excrement.
 - The location of any known populations, including grid references.
 - The date and time the sample was taken, how it was identified and by whom. If taken by non PHSI, an official sample would be required.
 - Current treatments/controls in place e.g. chemical treatments.
 - Details of the movement of people, equipment, machinery etc. to and from the infested area.
 - Cultural and working practices.
 - The name, address, Email and telephone number of the person who found the beetle.
- 5.16. This information should be included on the plant pest investigation template (see Appendix III of the Defra *Generic Contingency Plan for Plant and Bee Health in England*).
- 5.17. Further to information gathering, samples of other beetles should be taken to confirm the extent of the infestation e.g. in surrounding lots or fields. This initial survey will be used to determine if it is an isolated finding or an established outbreak.

- 5.18. When at a processor, any host plants (including any potato tubers) should be surveyed for signs of pest presence in the immediate vicinity (the premise in which the pest is found) of any outbreak if this occurs during the growing season and again during the following growing season (see 5.20-5.22). If the site is in a potato growing area, the two potato fields closest to the site and a field on the prevailing wind side should also be surveyed. This is a guide and may differ depending on the specific characteristics of the area. The waste disposal areas should also be inspected.
- 5.19. Finance for the surveys will depend on the individual circumstances of the outbreak, and will be subject to discussion, usually between Defra policy and the PHSI.

Sampling

- 5.20. Eggs, larvae and adults are distinctive (see Appendix A Morphology section). Leaves can therefore be visually inspected for these life stages. Signs of defoliation and characteristic black, sticky excrement can also be looked out for. Because the larvae and adults can be hidden among the foliage, they are best detected by shaking the host plant over a white sheet or tray.
- 5.21. When walking through a potato crop, two rows can be taken at a time, and any findings marked with flagged canes (ideally around the perimeter of any colonies).
- 5.22. Following the identification/capture of an egg, larva, pupa and/or adult, the samples should be sent for diagnosis as in point 5.1.

Diagnostic procedures

- 5.23. There are descriptions of eggs and larvae of *L. decemlineata* (e.g. Australian Government Department of Agriculture, 2013), but there are no keys, so morphological species determination of these life stages is not possible, and DNA sequencing would be needed.
- 5.24. For adults, there is a North American key to the genera in the tribe Doryphorini, including the genera most closely related to *Leptinotarsa*, as well as 31 of the 41 species in the genus *Leptinotarsa*, including *L. decemlineata* (Jacques, 1988).

Criteria for determining an outbreak

- 5.25. If *L. decemlineata* is detected at a location and is not confined to a particular consignment(s) then an outbreak should be declared. For example, in a potato field. However, if the finding is restricted to recently imported potatoes within a cold store or to other produce, then this would be classified as an interception.

Official action to be taken following the confirmation of an outbreak

- 5.26. The scale of the outbreak will determine the size and nature of the management team and action.

Communication

- 5.27. The IMT will assess the risks and communicate details to the IPPC, EU and EPPO, in accordance with ISPM 17: pest reporting (<https://www.ippc.int/en/publications/606/>), as well as to devolved authorities, within Government to Ministers, senior officials and other government departments and agencies (e.g., the Environment Agency) on a regular basis as appropriate; and to stakeholders (of which there is a prepared list).

Surveillance and demarcated zones

- 5.28. After an outbreak has been detected, a regulated area should be established that includes:
- An infested zone (i.e. the infested field or crops). This may also include field margins or uncropped areas if infestation is found on solanaceous weed species.
 - A buffer zone, which should extend out at least 1 km from the infested zone, but may extend out to 5 km depending on the characteristics of the outbreak.
- 5.29. Initial maps of outbreak sites should be produced by either the PHSI or the Risk and Horizon Scanning team, with ongoing mapping capabilities carried out by the latter. Grower contact details will be available from the AHDB buyers and grower levy payers list.
- 5.30. If it is considered possible that the pest has been spread to distant fields on machinery, these fields should also be surveyed. If resources are limited, priority should be given first to areas where there has been movement of large quantities of soil from the infested zone and secondary areas looked at as resources dictate.
- 5.31. All host plants in the buffer zone should be visually inspected and any suspect samples should be sent for diagnosis (5.20-5.22). This will include fields containing volunteer potato plants or solanaceous weeds.
- 5.32. The regulated area should be adjusted in response to further findings. If *L. decemlineata* is found within a field outside the infested zone, this should subsequently be designated as infested. If beetles are found within uncropped areas outside the infested zone, then any field directly adjacent to these areas should normally be designated as infested.

Decontamination procedures

- 5.33. Within the infested zone, all non-disposable material, equipment and machinery, should be thoroughly cleaned to remove the pest and any soil before movement to un-infested areas.
- 5.34. Any waste (plant or other potentially infested material) should be removed and destroyed (via deep burial, incineration or other appropriate methods, see 5.58-5.60).

Tracing forwards/backwards

- 5.35. Once other sites that are potentially infested by *L. decemlineata* have been identified, these should be inspected as per 5.15-5.22.
- 5.36. A pest alert to raise awareness of *L. decemlineata* and its symptoms should be published and distributed to potato growers, packers/processors and importers, and displayed at public locations, nearby to the outbreak site and other areas at risk.

Pest management procedures

Scenario 1: Outbreak in a potato crop – infested zone

- 5.37. The whole crop should be treated as soon as possible with a foliar insecticide. PHSI will advise on an appropriate treatment regime in consultation with the Defra Risk and Horizon Scanning team. Foliar insecticides will kill life stages above ground, but will have no effect on the larvae, pupae or adults within the soil.
- 5.38. If the crop is organic, pesticides will still have to be used if the situation demands it as the statutory requirement to eradicate *L. decemlineata* overrides any restrictions on the use of pesticides on organic farms.
- 5.39. Growers will be placed under notice to apply the recommended pesticides and make the applications using their own or contractor's equipment. Records of applications will be kept, including details of the amount of product and water used.
- 5.40. Bee advisors and local beekeepers should be contacted to inform them of any insecticide applications and their timing. Bee inspectors should be able to provide contact details.
- 5.41. Visual inspection should be used to assess the efficacy of insecticide treatments. This should not be carried out while the crop is still wet with insecticide.
- 5.42. When the level of adults is low (no adults detected), the potato haulm should be destroyed (using methods in point 5.57) to eliminate the food supply of *L. decemlineata* and thereby reduce their ability to survive and multiply.
- 5.43. If the level of adults is high (adults detected), the potato haulm should **not** be destroyed due to the high risk of adults dispersing to hosts in other fields/sites.
- 5.44. Even if the number of adults is low when the haulm is destroyed, there is still a risk of some adults (including newly emerging adults) spreading further afield. Several rows of crop should therefore be left at the edge of the field to act as a trap. These should be treated regularly (e.g. weekly) with insecticide and destroyed later in the season.
- 5.45. No tubers should be harvested due to the risk of pest dispersal with the tubers and any associated soil. Destruction of tubers in a way that does not risk further spread (e.g. exposure to frost) should be considered. The method of destruction will be decided by the IMT.
- 5.46. Host plants, such as solanaceous weeds, and host debris in the infested zone, and uncropped areas, field boundaries and hedgerows in the immediate vicinity of the infested zone should be destroyed by herbicide or mechanical means.

Scenario 2: Measures to be taken in the case of detection of infestation in tubers after harvest (e.g. in a growers potato store or during processing/packaging when the tubers can be traced back to a known crop or farm)

- 5.47. The following should be designated as infested:
 - The lot from which the sample was taken.

- The waste from the infested lot, such as soil and processed waste.
 - The equipment and other articles (e.g. machinery and packing material) which have been in contact with the lot.
 - The field where the lot was grown.
- 5.48. As in point 5.28, a buffer zone should be created that extends out to at least 1 km from the infested field(s) and store.
- 5.49. Fields or areas where potentially infested equipment, waste, and other articles, have been used should be surveyed, and any tubers harvested from these fields should be inspected.
- 5.50. There is the risk that other potato stocks may have become infested after harvest. This could have been through the transfer of soil in grading lines or storage boxes. Any tubers and associated containers in storage, and tubers linked to infested machinery and equipment, should therefore be classed as possibly infested and inspected for the presence of *L. decemlineata*.
- 5.51. Points 5.2 – 5.6 and 5.8 – 5.9 should be followed, but only destruction rather than re-export should be considered.

Crops growing within the buffer zone (1 km around the infested zone) in the year of the outbreak

- 5.52. If no infestation is found in the potato crops growing in the buffer zone following surveillance, then they should be treated with a programme of foliar insecticides under notice until harvest and monitored for any sign of *L. decemlineata*.
- 5.53. Growers of non-potato crops, but which contain host volunteers or weeds may be restricted under notice from undertaking any husbandry or harvesting of the crops until they have been inspected and found free of the beetle. These growers will be identified by the PHSI locally.
- 5.54. Potato tubers should also be inspected during and/or immediately after harvesting. This will also apply to non-potato crops in fields or other areas containing volunteer potato plants or solanaceous weeds.
- 5.55. All potato tubers that are harvested should be brushed and/or washed to remove soil and the pest prior to movement outside of the buffer zone. The potatoes should also only be marketed as ware potatoes and must not be used as farm saved seed.

Disposal plan

Infested growing host crops

- 5.56. The primary means of destruction of potato plants in a field is through herbicide application. Actives that can be used include glyphosate, diquat (e.g. Reglone), carfentrazone-ethyl (e.g. Spotlight plus) and glufosinate-ammonium (e.g. Harvest), but the Defra Risk and Horizon Scanning team should be consulted for appropriate treatments.

Infested harvested tubers/soil/plant debris

- 5.57. Adults, pupae and possibly larvae may be present with the harvested tubers, associated soil and plant debris. It is important that all of this material is disposed of safely so as to eliminate the pest. When deciding on the most appropriate method(s) of disposal, factors such as the likelihood of adults being present, the level of handling and transportation required, all need to be taken into account. For all methods, measures need to be taken to ensure that there is no risk of spread during transport and treatment or disposal. Material that can be moved safely should be destroyed by incineration at a licensed facility (if in small quantities) or deep burial. Disposal and/or destruction should be under the approval and supervision of the PHSI. If the material has to be moved off the premises, it should be contained within at least one sealed layer, and two layers if possible. Deep burial may be done at an approved landfill site, or on the site or nearby farm, but only in agreement with the local Environment Agency. Incineration must comply with appropriate waste management regulations, Environment Agency in England, Scottish Environment Protection Agency and Natural Resources Wales.
- 5.58. Other possible methods of destruction for potato tubers should be considered on a case by case basis and include heat sterilization, industrial processing (under official supervision), fermentation and composting, steaming and feeding to animals, and anaerobic digestion (minimum temperature of 55°C for 24 h without interruption with a hydraulic dwell time in the reactor of at least 1 day).
- 5.59. All objects designated as 'infested', such as equipment, machinery, storage facilities that may be contaminated with infested soil or potato material should be thoroughly cleansed to remove the pest and all soil. This should be carried out at the outbreak site or a site nearby in agreement with a Plant Health and Seeds Inspector. Any waste material generated should be bagged and sent for deep burial or incineration (as in 5.58).

Measures in subsequent seasons

Infested zone

- 5.60. No host or root crop should be planted for at least 2 years and until no volunteer plants have been found for 2 consecutive years (under notice). Trap crops planted in the year following the outbreak are exempt from this rule.
- 5.61. If trap crops are used, these should be planted early in the season to ensure they are present before the emergence of adult beetles in spring, and they should ideally be planted at the perimeter of the field (e.g. 6-24 m, but width dependent on the size of the field and situation) where beetles are more likely to overwinter. If possible, trap crops could be used in combination with a chemical attractant. These plants should be inspected regularly during spring and summer (e.g. fortnightly), and sprayed regularly (e.g. weekly) with an appropriate insecticide program as discussed with the Risk and Horizon Scanning team.
- 5.62. If practical, steam treatment and soil sterilants could be used prior to the emergence of adult *L. decemlineata*. The treated soil should subsequently be covered for three months to improve the efficacy of the soil treatment. For large fields, only areas of known infestation should be treated and covered.
- 5.63. Any volunteer plants should be destroyed early in the season following the outbreak. If the population of volunteer plants is low, they can be removed by hand, but if the population of

volunteer plants is high, they are best controlled by an application of an effective herbicide (see 5.57). Solanaceous weeds should also be controlled in a similar way.

- 5.64. 'Infested' fields may be maintained in permanent pasture with frequent close cutting or intensive grazing. This option has the advantage of providing effective control of potato volunteers and solanaceous weeds.
- 5.65. The frequency of inspections will be determined by the Incident Management Team.
- 5.66. Following 2 years without volunteer plants, only ware potatoes or a trap crop should be produced in the following season, with the crop and any harvested tubers inspected for *L. decemlineata*. If there are no finds of the pest following this, then either seed or ware potatoes can be produced on the field the following year.

Buffer zone (at least 1km around the infested zone)

- 5.67. Host crops should not be planted outdoors for at least 2 years (under notice). Volunteer plants and weeds should be destroyed. Following this period, only ware potatoes should be produced with the growing crop and harvested tubers inspected for *L. decemlineata*. If there are no finds of the pest, then either seed or ware potatoes can be produced on the field.
- 5.68. The frequency of inspections will be determined by the Incident Management Team.
- 5.69. Host crops planted indoors should be monitored, and if *L. decemlineata* found, the site (which may include outdoor fields) should be designated as infested and an appropriate programme of insecticides applied.

6. Criteria for declaring eradication / change of policy

- 6.1. A *Leptinotarsa decemlineata* outbreak can be declared eradicated (by the CPHO) only after at least three years during which time no *L. decemlineata* life stages have been found. These three years must include 2 consecutive years without volunteer plants and then a year with either a ware potato crop or trap potato crop.

7. Evaluation and review of the contingency plan

- 7.1. The Defra Contingency Plan for Plant Health in England requires that any pest specific plan is reviewed following an outbreak. This pest specific contingency plan should also be reviewed annually to take into account of changes in legislation, control procedures, pesticides, sampling and diagnosis methods, and any other relevant amendments.
- 7.2. Lessons should be identified during and after any *L. decemlineata* or other pest outbreak, including what went well and what did not. These should be included in any review of the contingency plan leading to continuous improvement of the plan and response to outbreaks.

8. Appendix A

Data sheet for *Leptinotarsa decemlineata*

Identity

PREFERRED SCIENTIFIC NAME	AUTHOR (taxonomic authority)
<i>Leptinotarsa decemlineata</i>	(Say, 1824)

CLASS: Insecta

ORDER: Coleoptera

SUBORDER: Polyphaga

SUPERFAMILY: Chrysomeloidea

FAMILY: Chrysomelidae

SUBFAMILY: Chrysomelinae

OTHER SCIENTIFIC NAMES

Chrysomela decemlineata Say

Doryphora decemlineata Say

Polygramma decemlineata Say

Leptinotarsa multitaeniata Stål

INTERNATIONALLY USED COMMON NAME/S & INTERNATIONAL LANGUAGE

Coloradobille (Danish)

Kartoffelbille (Danish)

Kartoffelkäfer (German)

Kolorado-Käfer (German)

Colorado beetle (English GB)

Colorado potato beetle (English US)

Ten-lined potato beetle (English GB)

Ten-striped sparmen (English GB)

Catarinita de la papa (Spanish)

Dorifora (Spanish)

Escarabajo de la patata (Spanish)

Escarabajo de las hojas de patata (Spanish)

Mayata de la papa (Spanish)

Tortuguilla (Spanish)
Susske sibsamini (Persian)
Koloradokuoriainen (Finish)
Chrysomèle de la pomme de terre (French)
Doryphore (French)
Doryphore de la pomme de terre (French)
Burgonyabogar (Hungarian)
Crisomela della patata (Italian)
Dorifora delle patate (Italian)
Scarabeo del Colorado (Italian)
Coloradokever (Dutch)
Koloradobille (Norwegian)
Stonka ziemniaczana (Polish)
Doriforo (Portuguese)
Escaravelho-da-batateira (Portuguese)
колорадский жук (Russian)
колорадский картофельный жук (Russian)
Koloradoskalbagge (Swedish)
Patetes bocegi (Turkish)

Notes on taxonomy and nomenclature

Leptinotarsa decemlineata is one of over 40 species in the genus *Leptinotarsa* (Arnett *et al.*, 2002). The genus is identified by the apical segment of the maxillary palpi (mouthparts) being shorter than the preceding segment, the mesosternum not being raised above the level of the prosternum, and the profemur of the male being simple (Jacques, 2000).

Biology and ecology

Life history

The adult beetles overwinter in the soil, usually in areas adjacent to potato fields, but also within the potato fields themselves. The depth at which the beetles overwinter in the soil varies depending on the climate, with beetles burrowing deeper (between 25 and 40 cm) in the cooler, northern European countries (CABI, 2016). Adults emerge over several weeks in spring and/or summer, and walk or fly to their host plants. They locate their host plants largely at random, though the odour of potato plants has been found to be attractive to the beetles; potato plants damaged by feeding were more attractive under laboratory conditions than undamaged plants (Bolter *et al.*, 1997; Landolt *et al.*, 1999). The adults are able to fly many kilometres in search of a host, including over large water bodies (Wikteliuss, 1981). Once they reach the potato plants, the adults feed and mating usually takes place within 5-10 days of emergence.

Eggs are laid in masses (of 10-30), usually on the underside of leaves, with each female laying up to 2000 eggs during their lifetime. Larvae use egg bursters (or oviruptors) on their meso and metathorax, and abdominal segments to hatch from the egg (within 4-14 days). They subsequently feed on the chorion (outer shell), before moving onto foliage within 24 hours. Larvae tend to hatch synchronously within egg masses and stay together until their first moult. Larvae have four instars,

with the level of feeding increasing between instars; 3, 5, 15 and 77% of consumption during larval development occurs in the first, second, third and fourth instars, respectively (Ferro *et al.*, 1985). Feeding generally takes place on the leaves, but larvae will feed on the petioles and stems if leaves are unavailable. Similarly, larvae and adults will feed on eggs of their own species under certain conditions (e.g. Shrod *et al.*, 1996). Larval development takes between 8 and 28 days depending on the temperature, and is largely not impacted by unfavourable weather conditions, though heavy rain and strong winds have been shown to cause mortality (CABI, 2016). Faithi *et al.* (2013), Kahrizeh *et al.* (2014) and Yaşar and Güngör (2005) have also found mortality, as well as fecundity and development time, to vary depending on the potato variety.

When fourth instar larvae mature, they burrow into the soil nearby to the plant they were just on and pupate. The pupal stage lasts from 8 to 20 days. Adults then emerge and feed on nearby plants. The number of generations varies between 1 and 4 depending on the climate; the number of day degrees required from the egg to the adult is approximately 543, with a development threshold of 10.90°C (Zhou *et al.*, 2010). Diapause is principally triggered by a short photoperiod, though it is also affected by temperature and host plant chemistry (Hsiao, 1988; Izzo *et al.*, 2014). The removal of host plants entirely can also trigger diapause, as is the case in North Carolina, where the beetle enters diapause following potato harvest in June despite the day lengths being at their longest (CABI, 2016). In mild climates, the beetles may undergo an additional aestival diapause during the summer. This is triggered by the deterioration of their host plants, and the beetles will come out of this diapause once the host plant condition has improved (CABI, 2016).

Hosts/crops affected

The most significant host of *L. decemlineata* is potato (*Solanum tuberosum*), though the beetle will feed on other cultivated crops, including tomato (*Solanum lycopersicum*) and aubergine (*Solanum melongena*), as well as wild solanaceous plants.

Table 1. Hosts of *Leptinotarsa decemlineata*.

Hosts	Type	Reference
<i>Brassica oleracea</i> (cabbage)	Minor	EPPO (2016)
<i>Cichorium endivia</i> (salad leaf)	Minor	EPPO (2016)
<i>Daucus carota</i> (wild carrot)	Minor	EPPO (2016)
<i>Lactuca sativa</i> (lettuce)	Minor	EPPO (2016)
<i>Petroselinum crispum</i> (parsley)	Minor	EPPO (2016)
<i>Nicotiana tabacum</i> (tobacco)	Minor	CABI (2016)
<i>Solanum lycopersicum</i> (tomato)	Minor	EPPO (2016)
<i>Solanum melongena</i> (aubergine)	Minor	EPPO (2016)
<i>Solanum tuberosum</i> (potato)	Major	EPPO (2016)
Solanaceous weeds	Wild/Weed	EPPO (2016)

Plant stage affected

Leptinotarsa decemlineata affects plants during growth and post-harvest.

Plant parts affected

Leaves, stems, vegetative organs, and tubers (occasional).

Symptoms/signs - description

Both the larvae and adults of *L. decemlineata* feed on foliage and strip the leaves from the edge inwards. Tubers, which are exposed at the soil surface, may also be eaten.

Additionally, larvae and adults produce black and sticky excrement, which is visible on the foliage.

Morphology

Egg: Orange/yellow in colour, elongate ovoid and approximately 1.2 mm in length. The female lays the eggs in upright clusters generally on the under surfaces of leaves where they are held in place by a special secretion produced by the female beetle. Image © Jyrki Tomminen (<https://creativecommons.org/licenses/by/2.0/legalcode>).



Larvae: First instar larvae are dark red and 1.5 mm in length. Each subsequent instar of which there are four in total is slightly paler than the last until by the 4th instar the mature larva is a paler orange red and up to 8mm long. The head and legs are consistently black in colour and there are two rows of black spots running along each side of the larvae. Typical chrysomelid larvae have a rather bloated, sack like and shiny body. Image © Stephan czuratis (<https://creativecommons.org/licenses/by-sa/2.5/legalcode>).



<p>Pupa: Yellowish in colour, naked (no pupal case is formed) and up to 1 cm in length. Pupation takes place in the soil, and so pupae are unlikely to be encountered. Image © Jyrki Tomminen (https://creativecommons.org/licenses/by/2.0/legalcode)</p>	
<p>Adults: Body oval, convex, 8.5 to 11.5 mm in length, males generally smaller. The elytra (wing cases) are smooth, shiny and yellow-orange in colour with each bearing 5 distinctive longitudinal black stripes. The head and pronotum (thorax) are orange in colour and are marked with a variable array of black spots. Legs are orange in colour. Image © Crown copyright</p>	

Similarities to other species/diseases/plant damages

A factsheet confirming the diagnostic characteristic of *L. decemlineata* and comparing this species with other insects frequently mistaken for it the UK is available

(https://secure.fera.defra.gov.uk/phiw/riskRegister/plant-health/documents/notifiable_pests/colorado-beetle-factsheet.pdf).

Detection and inspection methods

Leaves can be visually inspected for eggs, larvae and adult beetles. Also, signs of defoliation and characteristic black, sticky excrement can be looked out for. Because the larvae and adults can be hidden among the foliage, plants can be shaken over a white sheet or tray to dislodge any hidden specimens.

Distribution

Table 2. Global distribution of *Leptinotarsa decemlineata* (EPPO, 2016)

(P) present, (W) widespread, (L) localized, (O) occasionally present, (D) reported in the past, no longer present, (E) eradicated, (I) absent, intercepted only			
COUNTRY/REGION	DISTRIBUTION (see codes above)	COUNTRY/REGION	DISTRIBUTION (see codes above)

ASIA	P	CENTRAL AMERICA & THE CARIBBEAN	P
CHINA	L	CUBA	P
Xinjiang	P	GUATEMALA	P
IRAN	P	MEXICO	W
IRAQ	L	SOUTH AMERICA	Absent
JAPAN	I	EUROPE	P
KAZAKHSTAN	L	ALBANIA	P
KYRGYZSTAN	P	ANDORRA	P
TAJIKISTAN	L	ARMENIA	P
TURKMENISTAN	P	AUSTRIA	W
UZBEKISTAN	L	AZERBAIJAN	P
AFRICA	Absent	BELARUS	W
NORTH AMERICA	W	BELGIUM	W
CANADA	L	BOSNIA AND HERZEGOVINA	P
Alberta	P	BULGARIA	W
British Columbia	L	CROATIA	W
Manitoba	P	CZECH REPUBLIC	W
New Brunswick	P	DENMARK	E
Nova Scotia	P	ESTONIA	W
Ontario	P	FINLAND	L
Prince Edward Island	P	FRANCE	W
Québec	P	Mainland	P
Saskatchewan	P	Corse	P
USA	W	GEORGIA	P
Alabama	P	GERMANY	W
Arizona	P	GREECE	L
Arkansas	P	HUNGARY	W
Colorado	P	IRELAND	I
Connecticut	P	ITALY	P
Delaware	P	Mainland	W
Florida	P	Sicilia	P
Georgia	P	LATVIA	P
Idaho	P	LITHUANIA	W
Illinois	P	LUXEMBOURG	L
Indiana	P	MACEDONIA	P
Iowa	P	MOLDOVA	L
Kansas	P	NORWAY	I
Kentucky	P	POLAND	W
Louisiana	P	PORTUGAL	P
Maine	P	ROMANIA	W
Maryland	P	RUSSIA	L
Massachusetts	P	Central Russia	P
Michigan	P	Eastern Siberia	P
Minnesota	P	Far East	P
Mississippi	P	Southern Russia	P
Missouri	P	Western Siberia	P
Montana	P	SERBIA	W
Nebraska	P	SLOVAKIA	P
New Hampshire	P	SLOVENIA	W
New Jersey	P	SPAIN	P
New Mexico	P	Mainland	W
New York	P	Islas Baleares	L

North Carolina	P	SWEDEN	E
North Dakota	P	SWITZERLAND	P
Ohio	P	TURKEY	L
Oklahoma	P	UKRAINE	W
Oregon	P	UNITED KINGDOM	E
Pennsylvania	P	Channel Islands	E
Rhode Island	P	England	E
South Carolina	P	Northern Island	I
South Dakota	P	Scotland	I
Tennessee	P	OCEANIA	Absent
Texas	P		
Utah	P		
Virginia	P		
Washington	P		
West Virginia	P		
Wisconsin	P		
Wyoming	P		




Fig. 6. Distribution of *Leptinotarsa decemlineata* in Europe.

History of introduction/spread

Leptinotarsa decemlineata was first discovered in 1811 by Thomas Nuthall in the USA, and was later described by Thomas Say in 1824 from specimens collected in the Rocky Mountains on Buffalobur (*Solanum rostratum*) (Casagrande, 1985; Jacques, 1988). In 1859, likely as a result of

host range expansion, *L. decemlineata* was recorded damaging potato crops 100 miles west of Omaha, Nebraska, in the USA (Pope and Madge, 1984). The beetle gradually moved eastwards, crossing the Mississippi river in 1865, reaching Ohio in 1869 and Maine in 1872, before arriving at the Atlantic coast in 1874 (Casagrande, 1987; Jacques, 1988). It now occurs throughout most of the USA, with the exception of Alaska, California, Hawaii and Nevada. The beetle is also present in Canada, which it reached in 1901 (Ivanschik and Izhevsky, 1981), and Central America.

Leptinotarsa decemlineata was first recorded in Europe in 1875 in the UK, and soon after in Germany and Poland. Although these outbreaks were eradicated (Feytaud, 1950; Wegorek, 1955; Jacques, 1988), the beetle eventually became established in Europe in 1922, within Bordeaux, France, and subsequently spread across the rest of Europe. It is now found in Belarus, Belgium, Romania, Russia, Slovakia, Spain (1935 – year first reported), Estonia, the Czech Republic, Germany, Italy, Latvia, Libya, Lithuania, Luxembourg (1936), Moldova, the Netherlands, Switzerland (1937), Austria (1941), Portugal (1943), Poland (1946), Hungary (1947), Turkey (1949), Bulgaria (1958), Greece (1963) and several other European countries. The beetle has also moved further east, into Asia.

Since 1952, the UK has only experienced two outbreaks of *L. decemlineata*; one in 1976 and one in 1977. In September 1976, four egg batches and 135 adult beetles were found in a field of potatoes (184 m²) in Thanet, Kent. After the implementation of control measures, no further beetles were found after March of the following year. The second outbreak in July 1977 was also small and consisted of just 61 larvae in a 12 m² allotment in Hastings, Kent. Inspections carried out after treatment of the allotment found no further beetles.

While there have been no outbreaks since 1977, there have continued to be interceptions of the beetle. The beetles have predominantly been found “hitch-hiking” on leafy vegetables, salad leaves, fresh herbs and grain, though they have also been found on potato. It should be noted that the number of interceptions has fallen since 1996, and from 2009 to 2016, there have been less than 10 interceptions per year.

Phytosanitary status

Leptinotarsa decemlineata is a IB EU listed pest. Countries with protected zone status include Spain (Ibiza and Menorca), Ireland, Cyprus, Malta, Portugal (Azores and Madeira), UK, Sweden (Blekinge, Gotland, Halland, Kalmar, Skåne), and Finland (the districts of Åland, Turku, Uusimaa, Kymi, Häme, Pirkanmaa, Satakunta).

Leptinotarsa decemlineata is also on the EPPO A2 list and on many other quarantine organism lists (see table 3).

Table 3. Lists on which *Leptinotarsa decemlineata* has been included as a quarantine organism (EPPO, 2016).

Country/Organisation	List	Year of addition
East Africa	A1 list	2001
Southern Africa	A1 list	2001
Argentina	A1 list	1995

Brazil	A1 list	1992
Canada	A2 list	1995
Chile	A1 list	1992
Paraguay	A1 list	1992
Uruguay	A1 list	1992
China	A1 list	1988
Israel	Quarantine pest	2009
Jordan	Quarantine pest	2007
Uzbekistan	A2 list	2008
Azerbaijan	A2 list	2007
Belarus	Quarantine pest	1994
Norway	Quarantine pest	2012
New Zealand	Quarantine pest	2000
APPPC	A1 list	1988
CAN	A1 list	1992
COSAVE	A1 list	1992
CPPC	A1 list	1990
EPPO	A2 list	1975
EU	Annex 1/B	1992
IAPSC	A2 list	1989
OIRSA	A2 list	1992
PPPO	A2 list	1993

Means of movement and dispersal

Adults of *L. decemlineata* are not strong fliers and tend to remain in the area where they developed as larvae (Grafius, 1995; Hurst, 1975). However, they can travel considerable distances in search of new habitat when aided by the wind, sometimes by as much as 100 km or more (van Poeteren, 1939; Wiktelius, 1981). Adults are also capable of surviving for long periods on sea water (Feytaud, 1936; Vassiere, 1939); on more than one occasion, they have been found coming ashore onto Jersey, having dropped into the sea after flying from the Contentin peninsula in France, which is 14 miles away (Brokenshire *et al.*, 2012).

Because of their presence on leaves and stems, eggs, larvae and adults can be found in association with host plants for planting in trade (CABI, 2016). Larvae and adults, as well as pupae, can also be found with any associated soil, and adults can sometimes be found on potato tubers (CABI, 2016). Additionally, adults can be found to be “hitch-hiking” on non-host plants, transport and packaging, as well as with travellers and their baggage (CABI, 2016). The majority of interceptions in the UK in the last few years have been as a result of this “hitch-hiking” on leafy vegetables, salad leaves and other non-host material all from continental Europe (Fera interception records, 2016).

Control

Biological control

There are a large number of natural enemies that attack *L. decemlineata*, including insect predators and parasitoids, nematodes, fungi, bacteria and viruses (a full list is provided by CABI, 2016). However, these are generally ineffective, particularly when compared with chemical alternatives.

Transgenic plants, which express the gene for *Bacillus thuringiensis* subsp. *tenebrionis* Cry3A delta-endotoxin is highly toxic to *L. decemlineata*. It was approved for commercial use in the USA in 1995, but due to negative public perception, there is a reticence from growers to use this method. As resistant beetles will also be selected for by these plants, there is a need for resistance management programmes. These involve the use of refuge areas of plants which do not express the endotoxin, and so will allow non-resistant beetles to persist in the population.

Other biological control methods include antifeedants (e.g. Neem extract, Bezjak *et al.*, 2006; Igrc *et al.*, 2006; Zehnder and Warthen, 1988) and the use of RNA interference (Zhu *et al.*, 2011).

Chemical control

The use of insecticides is the most commonly used method for controlling *L. decemlineata*. Effective insecticides include chlorpyrifos (no longer approved for use on potatoes in the UK) and thiacloprid. Partly because of the dependency on chemical insecticides, *L. decemlineata* has developed resistance to a number of chemicals. According to the Arthropod Pesticide Resistance Database (2016), there have been 296 cases of resistance in 140 locations against 55 active ingredients. In Long Island, New York State, USA, for example, the beetle exhibited a 155 fold resistance to imidacloprid after just 3 seasons. It is therefore recommended that different insecticide classes are rotated with one another as part of a resistance management programme.

Field monitoring/economic threshold levels

The economic injury level (EIL) is the fewest number of insects (or level of defoliation) that will cause yield losses equivalent to the cost of insect management. In the case of *L. decemlineata*, defoliation levels of 20% during plant emergence, 30% during early to late bloom and 60% during late bloom to harvest have been developed by Zehnder *et al.* (1995). However, these thresholds vary depending on the location of the potato plants, current cost of treatment, as well as the statistical approach used (cf. Nault and Kennedy, 1998; Shields and Wyman, 1984). Economic injury levels have also been developed for insect densities; Mailloux *et al.* (1995), for example, report an EIL of 5.8 overwintered adults and 10 summer generation adults per plant, whereas larval densities vary from 0.14 – 0.82 per plant (Senanayake and Holliday, 1990) to 12 per stalk (Mailloux *et al.*, 1991).

Economic thresholds, conversely, are the points at which action should be taken to prevent a pest population reaching the EIL. In many cases where these have been used for *L. decemlineata*, yield losses were not too dissimilar to conventional management programmes. However, fewer insecticide applications were needed (Stewart and Dornan, 1990; Wright *et al.*, 1987; Zehnder *et al.*, 1995). In Virginia, USA, a computerized approach is used called the Potato-Insect Expert System (PIES). This takes into account the insect life stage, the potato growth stage, percentage defoliation and other factors, to decide whether an insecticide application is necessary (Vencill *et al.*, 1995). Likewise, a temperature-driven decision support system (SIMPLEP DSS) was designed for Europe. It is composed of two modules; the first (SIMLEP 1) forecasts the first occurrence of the hibernating beetles and the start of egg laying and the second (SIMLEP 3) predicts the occurrence of the

development stages. Following validation in a number of countries, this has proven to be very accurate, correctly predicting the periods of maximum egg laying and young larval presence in 90% of cases. SIMLEP 3 is used by Germany, Austria, and the western part of Poland.

In spite of these systems to reduce the number of insecticide sprays, the USA tends to adopt a preventative approach, such as the in furrow applications of imidicloprid (CABI, 2016).

Cultural controls and sanitary methods

The level of infestation by *L. decemlineata* can be reduced by as much as 90% by rotating the potato crop between fields (Lashomb and Ng, 1984; Wright, 1984). Ideally, a minimum distance of 0.5 km from the source of the beetle is required to fully benefit from a crop rotation strategy. This strategy can be maximized by including growers from a large area, allowing longer distance rotations and so reducing the ability of the beetle to locate a host (CABI, 2016).

Populations of the beetle can also be reduced by altering the planting date. Early planting means that the plants will mature before the second generation of beetles is produced, while late planting means that summer generation adults will emerge after the critical photoperiod in autumn and will not be able to produce a second generation (Weber and Ferro, 1993).

The majority of beetles tend to overwinter at the edges of potato fields (hundreds/m²), with only a few beetles per m² overwintering in the fields themselves (Hunt and Tan, 2000; Weber and Ferro, 1993). It is also known that beetles tend to disperse initially by walking rather than by flying. Trenches (which act as pitfall traps) and trap crops along the field margins can therefore be effective in preventing and/or slowing the entry of beetles into a field. Plastic lined trenches, for example, caught 95% of the beetles during field experiments (Boiteau and Osborn, 1999; Misener *et al.*, 1993). The efficacy of these traps can be improved further by using semiochemicals; Martel *et al.* (2005) showed that a blend of chemicals emitted by the host plants ((Z)-3 hexenyl acetate, \pm -linalool and methyl salicylate) increased the efficacy of pitfall traps and, in turn, reduced the level of pesticide needed in the crop.

Potato breeding lines for resistance have also been released (Plaisted *et al.*, 1992; Lorenzen and Balbyshev, 1997). These are largely based on resistance derived from *Solanum berthaultii*, which has glandular trichomes on the foliage and resistance derived from *Solanum chacoense*, which has high concentrations of leptine glycoalkaloids in the foliage (Sinden *et al.*, 1986; Sanford *et al.*, 1997; Yencho *et al.*, 1996, 2000).

Further control measures include mechanical collection, propane flamers, pneumatic thermal machines or bio-collectors and mulching (CABI, 2016).

Impacts

Economic impact

Leptinotarsa decemlineata is a destructive defoliator of potato plants worldwide (Hare, 1990), and is also responsible for spreading potato diseases, such as *Ralstonia solanacearum* and *Clavibacter michiganensis* subsp. *sepedonicus*.

In Michigan, with an area of 250,000 km², the combined cost of treatment and yield loss for potato was \$13.8 million in 1994, equating to 13.7% of the crop value (Grafius, 1997), whereas in China, total losses (not just including potato) of \$3.2 million were estimated in the infested area per year (Liu *et al.*, 2012). Potato yield losses of 50% in some EPPO countries, and tomato yield losses of 67% in a field test in Maryland, USA, have also been reported (EPPO, 1997; Schalk and Stoner, 1976).

Environmental impact

Leptinotarsa decemlineata is largely found on plants in the Solanaceae family, and primarily affects agricultural plants (e.g., potato) within this family; there are no indications that the beetle has any significant negative effects on wild plants in the environment. However, the beetle can have an indirect negative impact on arthropod biodiversity within agricultural fields because of the increased use of insecticides used to control it.

Social impact

Unlike *Phytophthora infestans* (potato blight), *L. decemlineata* has not caused enough damage to lead to a significant social impact.

9. References

- Australian Government Department of Agriculture** (2013). National Diagnostic Protocol for Colorado Potato Beetle, *Leptinotarsa decemlineata* [Online]. Available: <http://www.google.co.uk/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&ved=0ahUKEwjJlfNw7HOAhWB1cAKHXOeB2UQFgghMAA&url=http%3A%2F%2Fplantbiosecuritydiagnostics.net.au%2Fwordpress%2Fwp-content%2Fuploads%2F2015%2F03%2FNDP-22-Colorado-potato-beetle-Leptinotarsa-decemlineata-V1.2.pdf&usq=AFQjCNEfCc0V1AevWuahUxaDUnuVRTWI6w&bvm=bv.129391328,d.bGs>. Accessed: 08/08/2016.
- Arnett, R. H. Jr., Thomas, M. C., Skelley, P. E. and Frank, J. H.** (2002). American beetles. Volume 2. Polyphaga: Scarabaeoidea through Curculionoidea. CRC Press, Boca Raton. XIV + 861 pp.
- Arthropod Pesticide Resistance Database** (2016). *Leptinotarsa decemlineata* [Online]. Available: <http://www.pesticideresistance.org/display.php?page=species&arId=141>. Accessed: 08/08/2016.
- Bartlett, P. W.** (1980). Interception and eradication of Colorado beetle in England and Wales, 1958-1977. Bulletin, Organisation Europeenne et Mediterranee pour la Protection des Plantes. 10, 481-489.

- Bezjak, S., Igrc-Barcic, J. and Bazok, R.** (2006) Efficacy of botanical insecticides in Colorado Potato Beetle (*Leptinotarsa decemlineata* Say., Coleoptera@ Chrysomelidae) control (Učinkovitost boljnih insekticida u suzbijanju krumpirove zlatice (*Leptinotarsa decemlineata* Say, Coleoptera: Chrysomelidae).) *Fragmenta Phytomedica et Herbologica* . 29, 13-24.
- Boiteau, G. and Osborn, W. P. L.** (1999). Comparison of plastic-lined trenches and extruded plastic traps for controlling *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae). *Canadian Entomologist*. 131, 567-572.
- Bolter, C. J., Dicke, M., Loon, J. J. A., Visser, J. H. and Posthumus, M. A.** (1997). Attraction of Colorado potato beetle to herbivore-damaged plants during herbivory and after its termination. *Journal of Chemical Ecology*. 23, 1003-1023.
- Brokenshire, T., Meadows, S. and Genard, O.** (2012) An international campaign coordinated by EPPO to control Colorado Beetle in the Channel Islands. *EPPO Bulletin*. 42, 167-170.
- CABI** (2016). *Leptinotarsa decemlineata* (Colorado potato beetle) In: *Crop Protection Compendium*. Wallingford, UK: CAB International. www.cabi.org/cpc.
- Casagrande, R. A.** (1987). The Colorado potato beetle: 125 years of mismanagement. *Bulletin of the Entomological Society of America*. 33, 142-150.
- EPPO** (1997). *Quarantine Pests for Europe*. 2nd Edition. Ed. by Smith, I. M., McNamara, D. G., Scott, P. R. and Holderness, M. Wallingford, UK: CABI. 1425 pp.
- EPPO** (2016). EPPO Global Database *Leptinotarsa decemlineata* (LPTNDE) [Online]. Paris, France: European and Mediterranean Plant Protection Organization. Available: <http://www.eppo.int/DATABASES/pqr/pqr.htm>. Accessed: 08/08/2016.
- Faithi, S. A., Fakhr-Taha, Z. and Razmjou, J.** (2013). Life-history parameters of the Colorado potato beetle, *leptinotarsa decemlineata*, on seven commercial cultivars of potato, *Solanum tuberosum*. *Journal of Insect Science*. 13, 132.
- Ferro, D. N., Logan, J. A., Voss, R. H. and Elkinton, J. S.** (1985). Colorado potato beetle (Coleoptera: Chrysomelidae) temperature-dependent growth and feeding rates. *Environmental Entomology*. 14, 343-348.
- Feytaud, J.** (1936). Comment le doryphore envahit L'Europe. *Revue de Zoologie Agricole*. 35, 81 & 125.
- Feytaud, J.** (1950). Le Doryphore à la conquête de l'Europe. In: *Proceedings of the VIII International Congress of Entomology*, Sweden, Stockholm, pp. 643-646.
- Grafius, E.** (1995). Is local selection followed by dispersal a mechanism for rapid development of multiple insecticide resistance in the Colorado potato beetle? *American Entomologist*. 41, 104-109.
- Grafius, E.** (1997). Economic impact of insecticide resistance in the Colorado potato beetle (Coleoptera: Chrysomelidae) on the Michigan potato industry. *Journal of Economic Entomology*. 90, 1144-1151.

- Hare, J. D.** (1990). Ecology and management of the Colorado potato beetle. *Annual Review of Entomology*. 35, 81-100.
- Hunt, D. W. A. and Tan, C. S.** (2000). Overwintering densities and survival of the Colorado potato beetle (Coleoptera: Chrysomelidae) in and around tomato (Solanaceae) fields. *Canadian Entomologist*. 132, 103-105.
- Hurst, G. W.** (1975). Meteorology of the Colorado potato beetle. World Meteorological Organization Technical Note 137.
- Hsiao, T. H.** (1988). Host specificity, seasonality and bionomics of *Leptinotarsa* beetles. In: Jolivet, P., Petitpierre, E., Hsiao, T. H. eds. *Biology of Chrysomelidae*. Series entomologica. Volume 42. Dordrecht: Kluwer Academic Publishers. 581-599 pp.
- Igrc, J., Barcic, J., Dobrincic, R., Maceljski, M.** (1999). Effect of insecticides on the Colorado potato beetles resistant to OP, OC and P insecticides. *Anzeiger für Schädlingkunde*. 72, 76-80.
- Izzo, V. M., Armstrong, J., Hawthorne, D. and Chen, Y.** (2014). Time of the season: the effect of host photoperiodism on diapause induction in an insect herbivore, *Leptinotarsa decemlineata*. *Ecological Entomology*. 39, 75-82.
- Jacques, R. L.** (1988). The potato beetles: the genus *Leptinotarsa* in North America (Coleoptera, Chrysomelidae). *Flora and Fauna Handbook No. 3*. Leiden; New York: E. J. Brill. 144 pp.
- Lacey, L. A., Horton, D. R., Chauvin, R. L. and Stocker, J. M.** (1999). Comparative efficacy of *Beauveria bassiana*, *Bacillus thuringiensis*, and aldicarb for control of Colorado potato beetle in an irrigated desert agroecosystem and their effects on biodiversity. *Entomologia Experimentalis et Applicata*. 93, 189-200.
- Landolt, P. J., Tumlinson, J. H., Alborn, D. H.** (1999). Attraction of Colorado potato beetle (Coleoptera: Chrysomelidae) to damaged and chemically induced potato plants. *Environmental Entomology*. 28, 973-978.
- Lashomb, J. H. and Ng, Y. S.** (1984). Colonization by Colorado potato beetles, *Leptinotarsa decemlineata* (Say) (Coleoptera: Chrysomelidae), in rotated and nonrotated potato fields. *Environmental Entomology*. 13, 1352-1356.
- Liu, N., Li, Y. and Zhang, R.** (2012). Invasion of Colorado potato beetle, *Leptinotarsa decemlineata*, in China: dispersal, occurrence, and economic impact. *Entomologia Experimentalis et applicata*. 143, 207-217.
- Lorenzen, J. H. and balbyshev, N. F.** (1997). ND2858-1: a useful source of resistance to the Colorado potato beetle. *American Potato Journal*. 74, 331-335.
- Martel, J. W., Alford, A. R. and Dickens, J. C.** (2005). Synthetic host volatiles increase efficacy of trap cropping for management of Colorado potato beetle, *Leptinotarsa decemlineata* (Say). *Agricultural and Forest Entomology*, 7, 79-86.

- Mailloux, G., Binns, M. R. and Bostanian, N. J.** (1991). Density yield relationships and economic injury level model for the Colorado potato beetle larvae on potatoes. *Researches on Population Ecology*. 33, 101-113.
- Mailloux, G., Bostanian, N. J. and Binns, M. R.** (1995). Density yield relationships for Colorado potato beetle adults on potatoes. *Phytoparasitica*. 23, 101-118.
- Misener, G. C., Boiteau, G. and McMillan, L. P.** (1993). A plastic-lining trenching device for the control of Colorado potato beetle: Beetle Excluder. *American Potato Journal*. 70, 903-908.
- Nault, B. A. and Kennedy, G. G.** (1998). Limitations of using regression and mean separation analyses for describing the response of crop yield to defoliation: a case study of the Colorado potato beetle (Coleoptera: Chrysomelidae) on potato. *Journal of Economic Entomology*. 91, 7-20.
- Plaisted, R. L., Tingey, W. M. and Steffens, J. C.** (1992). The germplasm release of NYL 235-4, a clone with resistance to the Colorado potato beetle. *American Potato Journal*. 69, 843-846.
- Pope, R. D. and Madge, R. B.** (1984). The 'when' and 'why' of the Colorado potato beetle. *Antenna*. 8, 175-177.
- Sanford, L. L., Kobayashi, R. S., Deahl, K. L. and Sinden, S. L.** (1997). Diploid and tetraploid *Solanum chacoense* genotypes that synthesize leptine glycoalkaloids and deter feeding by Colorado potato beetle. *American Potato Journal*. 74, 15-21.
- Schalk, J. M. and Stoner, A. K.** (1976). Colorado potato beetle populations and their effect on tomato yield in Maryland. *HortScience*. 11, 213-214.
- Schrod, J., Basedow, T., Langenbruch, G. A.** (1996). Studies on bionomics and biological control of the Colorado potato beetle (*Leptinotarsa decemlineata* Say, Col., Chrysomelidae) at two sites in southern Hesse (FRG). *Journal of Applied Entomology*. 120, 619-626.
- Senanayake, D. G. and Holliday, N. J.** (1989). Economic injury levels for Colorado potato beetle (Coleoptera: Chrysomelidae) on 'Norland' potatoes in Manitoba. *Journal of Economic Entomology*. 83, 2058-2064.
- Shields, E. J. and Wyman, J. A.** (1984). Effect of defoliation at specific growth stages on potato yields. *Journal of Economic Entomology*. 77, 1194-1199.
- Sinden, S. L., Sanford, L. L. and Deahl** (1986). Segregation of leptine glycoalkaloids in *Solanum chacoense* Bitter. *Journal of Agricultural and Food Chemistry*. 34, 372-377.
- Stewart, J. G. and Dornan, A. P.** (1990). Comparison of three management schemes for Colorado potato beetle on early-season potatoes in Prince Edward Island. *Phytoprotection*. 71, 121-127.
- Van Poeteren, N.** (1939). Die Entwicklung der Karoffelkafer. Frage in Niederlanden. In *Verh. VII Intern. Kongr. Entomol., Berlin, 1938*, 2701-2703.

- Vassiere, M. P.** (1939). Au sujet de la dispersion du doryphore. Bulletin de la Societe Entomologique de France. 44, 178.
- Vencill, A. M., Zehnder, G. W. and Heatwole, C. D.** (1995). Potato insect expert system: computerized approach to insecticide management for Colorado potato beetle (Coleoptera: Chrysomelidae). Journal of Economic Entomology. 88, 944-954.
- Weber, D. C. and Ferro, D. N.** (1993). Distribution of overwintering Colorado potato beetle in and near Massachusetts potato fields. Entomologia Experimentalis et Applicata. 66, 191-196.
- Wegorek, W.** (1955). Investigation on spring migration of the Colorado beetle (*Leptinotarsa decemlineata* Say) and possibilities of combating the insect. Ekologia Polska Seria A. 3, 217-271.
- Wikteliu, S.** (1981). Wind dispersal of insects. Grana. 20, 205-207.
- Wright, R. J.** (1984). Evaluation of crop rotation for control of Colorado potato beetles (Coleoptera: Chrydomelidae) in commercial potato fields on Long Island. Journal of Economic Entomology. 77, 1254-1259.
- Wright, R. J., Kain, D. P., Moyer, D. D.** (1987). Development and implementation of an extension IPM program for Long Island. Bulletin of the Entomological Society of America. 33, 239-245.
- Yaşar, B. and Güngör, M. A.** (2005). Determination of life table and biology of Colorado potato beetle, *Leptinotarsa decemlineata* Say (Coleoptera: Chrysomelidae), feeding on five different potato varieties in Turkey. Applied Entomology and Zoology. 40, 589-596.
- Yencho, G. C., Bonierbale, M. W., Tingey, W. M., Plaisted, R. L. and Tanksley, S. D.** (1996). Molecular markers locate genes for resistance to the Colorado potato beetle, *leptinotarsa decemlineata*, in hybrid *Solanum tuberosum* x *S. berthaultii* potato progenies. Entomologia Experimentalis et Applicata. 81, 141-154.
- Yencho, G. C., Kowalski, S. P., Kennedy, G. G. and Sanford, L. L.** (2000). Segregation of leptine glycoalkaloids and resistance to Colorado potato beetle (*Leptinotarsa decemlineata* (Say)) in F2 *Solanum tuberosum* (4x) x *S. chacoense* (4x) Potato progenies. American Journal of Potato Research. 77, 167-178.
- Zehnder, G. and Warthen, J. D.** (1988). Feeding inhibition and mortality effects of neem-seed extract on the Colorado potato beetle (Coleoptera: Chrysomelidae). Journal of Economic Entomology. 81, 1040-1044.
- Zehnder, G., Vencill, A. M. and Speese, J.** (1995). Action thresholds based on plant defoliation for management of Colorado potato beetle (Coleoptera: Chrysomelidae). Journal of Economic Entomology. 88, 155-161.
- Zhu, F., Xu, J. J., Palli, R., Ferguson, J. and Palli, S. R.** (2011). Ingested RNA interference for managing the populations of the Colorado potato beetle, *Leptinotarsa decemlineata*. Pest Management Science. 67, 175-182.

10. Authors and reviewers

Authors:

Matthew Everatt (Defra)

Reviewers:

Dominic Eyre (Defra)

Ian Sanders (APHA)

Jennifer Baron (Defra)

Joe C. Ostojá-Starzewski (Fera Science Ltd.)