

Rapid Pest Risk Analysis (PRA) for:

Phytophthora pluvialis

April 2022

Summary and conclusions of the rapid PRA

This rapid PRA has been undertaken following a finding of the Oomycete pathogen *Phytophthora pluvialis* in Cornwall in September 2021. Subsequently, additional findings have been made elsewhere in England, also in Scotland and Wales on western hemlock and Douglas fir but not in Northern Ireland. Prior to the UK detections *P. pluvialis* was only known to be present in New Zealand and the Pacific Northwest (PNW) of the USA. In New Zealand it causes a disease known as red needle cast, primarily on radiata pine and Douglas fir, causing premature needle loss and loss of increment growth but trees recover. In the PNW it causes very minor damage on Douglas fir or tanoak and is considered part of the native forest mycota. This PRA shows:

Risk of entry

P. pluvialis is already present in the United Kingdom, the PRA area, but with a limited distribution across the UK and is the subject of official control. With the current plant health measures that are in place the likelihood of further entry via various pathways is assessed. Entry via pathway (1) traded **plants for planting** is considered as **very unlikely** (medium confidence); pathway (2) non-traded plants as **unlikely** (low confidence); pathway (3) timber/WPM as **very unlikely** (high confidence); pathway (4) cut conifer foliage as **very unlikely** (medium confidence); pathway (5) seeds as **unlikely** (low confidence); or pathway (6) soil and contaminated forestry machinery as **very unlikely** (low confidence).

Risk of establishment

Establishment under protection is considered **very unlikely** (high confidence). *P. pluvialis* has already established outdoors in some areas of England, Wales, and Scotland (high confidence), with natural spread via aerial spore dispersal and in water courses likely to

occur at **moderate** pace (medium confidence). In contrast, spread in trade via infected plants is considered likely to occur **quickly** (medium confidence), but **slowly** (low confidence) through the movement of wood/logs, and only **very slowly** (high confidence) via cut foliage. Lack of data on the potential host range of *P. pluvialis* and likelihood of sporulation on different plant parts affects confidence in these ratings.

Economic, environmental, and social impact

Based on uncertainty about host range, and the contrast in symptoms on affected trees in the UK compared with those in the PNW or New Zealand, the potential economic impact of *P. pluvialis* is rated **large** (low confidence), whilst both environmental and social impacts are assessed as **medium** (low confidence), with low confidence rating due to lack of data.

Endangered area

Susceptible hosts and a suitable climate for disease are present throughout much of the PRA area. The most favourable climatic conditions are likely to be throughout western UK based on comparisons with New Zealand and PNW where *P. pluvialis* is most active.

Risk management options

Eradication efforts are judged unlikely to succeed based on the number of findings already made. Short-term options for consideration include: (1) Containment of outbreak areas and host removal on only the most severely affected sites. (2) Controlled timber movement from affected sites. (3) Monitoring tree recovery on more lightly affected sites and what triggers serious disease episodes to understand the impacts of *P. pluvialis* over time.

Medium to longer-term management options include: (1) Use of silvicultural treatments to reduce the likelihood of severe disease expression. (2) Deployment of disease suppression compounds which reduce disease prevalence through targeted aerial application. (3) Breeding for resistance as a long-term option for high value hosts.

Key uncertainties/topics that would benefit from further investigation

Section of PRA	Further work needed to improve the PRA
Hosts (host range)	 Host range testing for susceptibility to <i>P. pluvialis</i> including UK grown <i>Pinus</i> species and other conifer species. Testing non-conifer species including species within the Fagaceae and some ornamental species.
	 Assessing the recovery potential of infected trees.
Pathogen behaviour	 Determining the rate of spread of the pathogen through epidemiological modelling, and climate matching to identify parts of the UK most at risk and inform future planting plans. Potential for persistence in soil and plant parts. Sporulation potential on UK grown hosts and optimum/minimum conditions for sporulation.
Pathogen genetics	Genotype comparisons of UK, NZ and USA <i>P. pluvialis</i> populations.
	• Potential for hybridisation with other Phytophthora species

Impact	 Loss of timber value and other public goods and services provided by affected trees
	 Potential for recovery associated with differing disease levels
Management	Control options for the pathogen in plantation and nursery situations.

Images of the pest

Photo 1



Source/ copyright owner: Forest Research

Source/ copyright owner: Forest Research



Is there a need for a detailed PRA or for a more detailed analysis of particular sections of the PRA? If yes, select the PRA area (UK or EU) and the PRA scheme (UK or EPPO) to be used.

No		
Yes	PRA area: UK or EU	PRA scheme: UK or EPPO

Given the information assembled within the time scale required, is statutory action considered appropriate / justified?

[The text below is a recommendation by the risk analyst which requires approval by PHRG]



Phytophthora pluvialis is newly identified in the UK and has already established outdoors in some areas of England, Wales, and Scotland. Due to the developing situation with this pest no final judgement has been made at present as to whether it meets the criteria for a Quarantine organism. Management options are continuing and evidence from all options plus ongoing research findings will be reviewed after 6-12 months to update disease management and assess the UK situation.

Stage 1: Initiation

1. What is the name of the pest?

Name:	Phytophthora pluvialis Reeser, Sutton & Hansen
Synonyms:	None
Taxonomy:	Kingdom – Chromista: Phyllum - Oomycota; Order - Peronosporales; Family - Peronosporacae; Genus - <i>Phytophthora</i>
Common name:	The pest does not have a common name, but the disease that it causes is referred to as red needle cast (RNC) in New Zealand
Etymology:	Specific ephithet 'pluvilalis' refers to the rain-associated canopy drip in tanwood-Douglas fir forests in western Oregon from which the first isolates of <i>P. pluvialis</i> were recovered from.

Special notes on taxonomy

The Internal transcribed spacer ITS-based phylogenetic tree for *Phytophthora* places *P. pluvialis* in Clade 3a, with closest relatives comprising *P. pseudosyringae*, *P. nemorosa*, *P. ilicis* and *P. psychrophila* (Abad et al. 2019).

2. What initiated this rapid PRA?

Severe decline was observed in a mature stand of western hemlock (*Tsuga heterophylla*) in southwest England (UK) in late August 2021, during the Forestry Commission's annual aerial surveillance for *Phytophthora ramorum*. Symptomatic material was collected and sent for analysis to the Tree Health Diagnostic and Advisory Service at Forest Research, UK. Analysis of material, including isolation onto Phytophthora-selective medium, yielded a species of *Phytophthora*. Based on ITS and coxII sequences the species was identified as *P. pluvialis* (Pérez-Sierra et al. 2022a); identification of *P. pluvialis* was also confirmed by real-time PCR using the protocol of McDougal et al. (2021). Previously *P. pluvialis* has only been known from New Zealand and the Pacific Northwest (PNW) states of the USA. It causes minor effects on Douglas fir (*Pseudotsuga menziesii*) and tanoak (*Notholithocarpus densiflorus*) in Oregon and wider PNW (Reeser et al. 2013; Hansen et al. 2015) and a more damaging needle disease on radiata pine (*Pinus radiata*) in New Zealand which Dick et al. (2014) named red needle cast (RNC). The pest was recognised as a potential risk to forestry species in the UK and had been added to the Plant Health Risk Register in 2014, but the recent findings in southwest England now require a PRA.

3. What is the PRA area?

The PRA area is the whole of the United Kingdom of Great Britain and Northern Ireland.

Stage 2: Risk Assessment

4. What is the pest's status in the plant health legislation, and in the lists of EPPO¹?

The pest is not listed in the EU² or GB³ plant health legislation and is not recommended for regulation as a quarantine pest by EPPO or on the EPPO Alert List. However, the UK Plant Health Risk Group concluded that *P. pluvialis* meets the criteria to be classified, at least initially, as a GB quarantine pest for regulatory purposes. Official measures currently in place include the prohibition of movement of any wood, isolated bark, and trees (including live trees, felled, or fallen trees, fruit, seeds, leaves or foliage) of the known host genera (*Tsuga, Pseudotsuga, Pinus* and *Notholithocarpus*) that originate within demarcated outbreak areas in GB⁴.

5. What is the pest's current geographical distribution?

The EPPO Global database shows the current known distribution of the pest (see Figure 1 reproduced from EPPO) with further details in Table 1 below.



Figure 1: Distribution of Phytophthora pluvialis (EPPO 2022)

¹ <u>https://www.eppo.int/ACTIVITIES/quarantine_activities</u>

² http://data.europa.eu/eli/reg_impl/2019/2072/oj

³ https://www.legislation.gov.uk/uksi/2020/1527/contents/made

⁴ <u>https://www.gov.uk/guidance/phytophthora-pluvialis</u>

Table 1: Distribution of Phytophthora pluvialis (EPPO 2022 and unpublished data)						
North America:	Present: California, Oregon, Washington, USA. Extent of the distribution may be under-estimated due to masking of symptoms by other pathogens.					
Central America:	Absent					
South America:	Absent					
Europe:	Present: UK (England, Scotland and Wales) but with restricted distribution.					
Africa:	Absent					
Asia:	Absent					
Oceania:	Present: New Zealand (North and South islands)					

6. Is the pest established or transient, or suspected to be established/transient in the UK/PRA Area?

Yes. Findings of the pest have been made in England, Scotland, and Wales. It has been found infecting established trees in managed forest/woodland environments. There have not been any findings in nurseries. Currently the pest is considered to have a limited distribution and is the subject of official control. Extensive surveillance by the Forestry Commission (FC), Scottish Forestry (SF), Natural Resources Wales (NRW), the Animal & Plant Health Agency (APHA) and Department of Agriculture, Environment and Rural Affairs (DAERA) in Northern Ireland is being undertaken to determine the distribution of *P. pluvialis* in the PRA area.

Summary of current situation

Between September 2021 and the end of April 2022, *P. pluvialis* was detected in the wider environment at 13 sites in England (Cornwall, Cumbria, Devon, Gloucestershire, Shropshire, and Surrey), 17 in Wales and five in Scotland (Highland and Argyll). At most of these sites western hemlock is the only known host, but both western hemlock and Douglas fir are affected at a few of the commercial forestry plantations. There have been no findings in the wider environment in Northern Ireland.

7. What are the pest's natural and experimental host plants; of these, which are of economic and/or environmental importance in the UK/PRA area?

Host plants

All known hosts of *P. pluvialis* are within the family Pinaceae, except for *Notholithocarpus densiflorus* which is in the Fagaceae. Symptoms incited by *P. pluvialis* vary with host and are summarised in Table 2, but much of the information is incomplete for minor hosts. Details of symptoms are presented in host and country combinations, as the same disease symptoms are not always apparent in each host for pest outbreaks reported from different countries.

Table 2: Naturally infected hosts of Phytophthora pluvialis							
Host species	Common name	Disease type	References				
Notholithocarpus densiflorus	Tanoak	Twig and stem cankers (USA)	Reeser et al. 2013				
Pinus radiata	Radiata pine	Needle cast (NZ)	Dick et al. 2014				
Pinus patula	Patula pine	Needle cast (NZ)	Scott et al. 2019				
Pinus pinea	Stone pine	Needle cast (NZ)	Scion 2022 (unpublished report)				
Pinus strobus	Weymouth pine	Needle cast (NZ)	Scott et al. 2019				
		Needle cast (NZ)	Gómez-Gallego et al. 2017				
Pseudotsuga mensiezii	Douglas fir	Needle cast and twig lesions (USA)	Hansen et al. 2015				
		Stem cankers and needle cast (UK)	Records FR THDAS 2021; Pérez-Sierra et al. 2022b				
Tsuga heterophylla	Western hemlock	Needle cast (NZ)	Scion 2022 (unpublished report)				

	Twig and stem cankers, needle cast (UK)	Pérez-Sierra et al. 2022a
	needie cast (UK)	2022a

New Zealand

Radiata pine and Douglas fir are the primary hosts of *P. pluvialis* in New Zealand, with single or occasional reports of the agent infecting *Pinus pinea* (stone pine), *P. patula* (patula pine), *P. strobus* (Weymouth pine) and western hemlock. With all hosts, infection appears almost exclusively limited to needles and invariably leads to premature needle loss.

Early symptoms on radiata pine consist of olive-coloured lesions on needles that contain black, resinous bands. The lesions quickly turn khaki-coloured with entire needles then taking on a yellow-brown or red hue before being cast, hence the common name of red needle cast (RNC) coined by Dick et al. (2014). A thick carpet of prematurely shed needles on the forest floor can be a sign of heavy infection. Apart from causing symptoms on needles, *P. pluvialis* has also been isolated from the rhizosphere of diseased radiata pine grown in bare-rooted nursery beds, but not from naturally infected roots (Scott et al. 2019). However, following artificial inoculation, *P. pluvialis* has been shown to have the potential to infect and limit the growth of fine roots of radiata pine at apparently sub-lethal levels (Scott et al. 2019).

Symptoms on Douglas fir are similar to those found on radiata pine: needles have olivecoloured lesions with black resinous bands, so needles take on an overall mottled, chlorotic appearance (Gómez-Gallego et al. 2017). They are dislodged readily so suitable test material for diagnosis can be lost rapidly (Gardiner et al. 2020). Experimental infection with zoospores of *P. pluvialis* reproduces the symptoms in Douglas fir, although the presence of another pathogen *Nothophaeocryptopus* (*Phaeocryptopus*) gaeumannii (Swiss needle cast) produces similar symptoms and may interact with *P. pluvialis* or mask its presence (Gómez-Gallego et al. 2017).

For minor hosts occasionally infected by *P. pluvialis* (patula pine, stone pine, Weymouth pine and western hemlock), there is little detail about the symptoms but banding or necrotic spotting on needles and premature needle loss is mentioned, although the pattern of needle cast differs from that observed on radiata pine (Scion 2022). Disease outbreaks on these minor hosts usually only occur when they are growing adjacent to radiata pine and Douglas fir stands already heavily affected by RNC. This suggests that minor hosts only become infected by *P. pluvialis* when exposed to high inoculum loads and so are less susceptible to the pathogen.

Pacific Northwest (PNW), USA

Douglas fir and tanoak are the only known hosts of *P. pluvialis* in the USA. First reports of *P. pluvialis* came in the absence of any visibly diseased hosts after its recovery from streams, soil samples and canopy drip in mixed tanoak-Douglas fir forest in Oregon (Reeser et al. 2013). Surveys then detected *P. pluvialis* on rare occasions causing twig

and stem cankers on tanoak, although it was found to be only weakly pathogenic when inoculated into tanoak stems (Reeser et al. 2015).

Findings on Douglas fir were made by Hansen et al. (2015) and it is now thought that *P. pluvialis* is widespread but a usually inconspicuous foliar pathogen when associated with this host (Hansen et al. 2017). Reported symptoms are mainly chlorotic needles that are shed readily. Two-year old seedlings exposed to natural inoculum levels in the forest have been found to develop twig symptoms that included tip dieback and stem lesions extending from bud scars, in addition to premature needle loss and irregular, mottled needle chlorosis. The same type of twig symptoms could not be seen on overstorey trees (Hansen et al. 2015). Similar symptoms have been induced on experimentally inoculated Douglas fir seedlings. However, in winter 2014-2015, specific symptoms associated with *P. pluvialis* consisting of dramatic chlorosis/ reddening of needles were observed on Douglas fir trees of all ages in many locations in the central coast range of Oregon. By the following year symptoms had declined and the trees recovered (Hansen et al. 2017).

Great Britain

On the most frequently affected host, western hemlock, *P. pluvialis* causes dieback, premature needle drops, and branch and stem cankers on mature trees. Young, naturally regenerated hemlock in the understorey of affected mature trees can also show high levels of dieback and mortality (Pérez-Sierra et al. 2022a). Branch and stem cankers exude copious resin and often have a blackened and cracked surface. When the outer bark of the cankers is removed to expose necrotic lesions in the phloem, resin pockets are usually visible below the lesions and sapwood (Pérez-Sierra et al. 2022a). Just as with radiata pine, it is usually the lower branches of trees that are affected first, with browning needles and multiple cankers on twigs and branches of all sizes. Observations also indicate that cankers can occur at the root collar and on major roots (A. Pérez-Sierra, unpublished data), and fine root systems on understorey hemlock may also be impaired (C.M. Brasier, unpublished data).

With Douglas fir, symptoms are less severe, but include chlorotic and prematurely cast needles, and stem cankers with underlying resin pockets (Pérez-Sierra et al. 2022b). However, as observed elsewhere with this host, *N. gaeumannii* (Swiss needle cast) infection of needles may also be present and cause similar symptoms to *P. pluvialis* on needles.

Apart from satisfying Koch's Postulates on western hemlock (Pérez-Sierra et al. 2022a), tests to challenge other conifer and broadleaf hosts with *P. pluvialis* to assess the potential host range have yet to be undertaken. However, during monitoring using detached sections of foliage or leaves ('baits') to detect the presence of *P. pluvialis* in water courses, Douglas fir and western hemlock shoots were colonised by natural available inoculum of the pathogen. Furthermore, on one occasion foliage of Koyama spruce (*Picea koyamae*) and on another occasion foliage of Sitka spruce (*P. sitchensis*) acted as a bait for *P. pluvialis* suggesting some susceptibility, although systematic testing of hosts from a range of genera has yet to be undertaken.

Economic and environmental importance of host plants

Tanoak is occasionally planted in the UK as an ornamental tree, with very few suppliers of the species listed in Britain⁵. Radiata pine is more common, but forestry plantings are mainly limited to provenance trials (see <u>http://www.forestry.gov.uk/fr/INFD-8CVE4W</u>). The Botanical Society of Britain and Ireland (BSBI) Atlas (Figure 2) shows the distribution of the species as mainly limited to southwest England and the coastal regions of Wales and Southern England, with very sporadic records in the central belt right up to Northeast Scotland. Douglas fir is considered a principal tree species, widely used for forestry and a popular and proven option to diversify production forests in line with the UK forestry standard⁶. It is found throughout much of the British Isles (Figure 2). Planted areas comprise just under 60,000 ha across Britain (Table 3), and amount to about 2% of the total conifer stock in Britain⁷. In contrast, western hemlock is considered a minor forestry species (less than 0.5% of the total conifer stock) and has a more limited distribution (Table 3) with much of the 9,000 ha of stock concentrated in southern and southwest England (Figure 2; Harmer et al. 2011). None of the known hosts are native to Britain which reduces their value to the environment.

Table 3: Planted areas of Douglas fir and western hemlock in Britain (ForestryStatistics 2021)

Conifer species	England	Scotland	Wales	Great Britain
Douglas fir	24,751 ha	24,565 ha	9,690 ha	59,006 ha
Western hemlock	5,706 ha	1,849 ha	1,512 ha	9,067 ha
Total	30,457 ha	26,414 ha	11,202 ha	68,073 ha

⁵ RHS Plantfinder <u>https://www.rhs.org.uk/plants/nurseries-search-result?query=10348</u>

⁶ https://www.gov.uk/government/publications/the-uk-forestry-standard

⁷ Forestry Statistics 2021: https://www.forestresearch.gov.uk/tools-and-resources/statistics/data-downloads/



Figure 2: Distribution of radiata pine, Douglas fir and western hemlock based on Botanical Society of Britain and Ireland (BSBI) maps showing presence in 10 km squares in Great Britain and Ireland <u>https://bsbi.org/maps</u>

8. Summary of pest biology and/or lifecycle

Phytophthora pluvialis is a homothallic species, meaning it can undergo sexual reproduction and as a result produce oogonia (sexual spores) in single-strain culture. *Phytophthora* oogonia are often considered to be persistent spores, able to survive for weeks or months under suboptimal conditions (Erwin & Ribeiro 1996; Boevink et al. 2020). The more ephemeral spore type, sporangia - which release free-swimming zoospores, are formed through asexual reproduction. The sporangia are semi-papillate and partially caducous (easily shed) (Reeser et al. 2013) indicating their potential for aerial dispersal, in rain splash and fog.

Phytophthora pluvialis causes a polycyclic disease, and studies of the pathogen in New Zealand have shown there are several infection cycles per year that are closely associated with rainfall (Gómez-Gallego et al. 2019a; Williams & Hansen 2018). Spores (presumed to mostly be sporangia/ zoospores) of *P. pluvialis* are released from infected foliage during the cooler period of the year between autumn and spring (Fraser et al. 2020) and in the presence of free-water on needle surfaces, infection can occur within 18 hours, with proliferation in plant tissue producing masses of sporangia which start to protrude from needle stomata a few days later (Gómez-Gallego et al. 2019a). Providing there are conducive weather conditions, reinfection cycles will occur every 4-6 days. In New Zealand the first symptoms of RNC usually appear in autumn or winter on the lower branches of affected trees, but under conducive conditions the disease can spread up the crown and to neighbouring trees. Where symptoms are limited to the lower branches, and sporulation occurs on infected foliage on these branches, aerial dispersal of *P. pluvialis* may remain largely within the understorey layer and the potential for longer distance dispersal above the tree canopy be much more limited.

Infected needles are cast by early to mid-spring in the following year (Dick et al. 2014; Fraser et al. 2020). It is unclear once the needles are cast and in the litter layer, if the inoculum they contain plays any further role in the disease cycle. Potentially, inoculum of oospores and asexual resting structures (e.g. encysted zoospores, stromata, hyphal swellings/aggregations) of *P. pluvialis* could be present in the litter layer and rhizosphere, and act as a reservoir of genetic diversity and inoculum for reinfection. Typically, oospores generally are resistant to extreme environmental conditions, and can survive for several years in soil or host plant material (Erwin & Ribeiro 1996). However, oospores or other resilient structures have rarely been observed in plant material (Hood et al. 2014; Williams & Hansen 2018) so the mode of survival by P. pluvialis through warmer/drier summer months has yet to be elucidated. Under favourable conditions for P. pluvialis, affected trees that were completely green at the start of autumn can be defoliated almost completely by the following spring, but the new growth is seldom affected (Dick et al. 2014). Therefore, the disease affects photosynthetic capability and hence growth, but does not cause tree mortality at least in New Zealand (Ganley et al. 2014). In the USA, P. pluvialis is considered an inconspicuous foliar pathogen which is probably native and causes relatively little damage to hosts (Reeser et al. 2013; Hansen et al. 2017).

9. What pathways provide opportunities for the pest to enter and transfer to a suitable host and what is the likelihood of entering the UK/PRA area?

Phytophthora pluvialis has been confirmed at a limited number of sites (England, Scotland, and Wales) across the PRA area. The most northerly finding is in Ross-shire⁸, furthest west in Cornwall, and most easterly in Surrey. In the same way that it is now considered that *P. pluvialis* was introduced into New Zealand several years before RNC became noticeable on radiata pine in 2008 (Dick et al. 2014), it is also likely that the pathogen has been in the UK for some time although visible signs of disease may only have emerged in 2021, possibly in response to a sequence of years with favourable weather conditions (Defra, unpublished data 2022). For many of the affected sites, symptoms observed in autumn through to spring in 2021-22 may not have been noticed or considered concerning were it not for the specific surveys and interest triggered by the finding of *P. pluvialis* on a number of heavily affected trees at a site in the southwest of England. Several possible pathways exist for entry into regions of the PRA area, with some likely to be direct/principal pathways (plants, wood) with others probably less significant or indirect pathways (seeds, foliage, soil, and machinery). The risk from each pathway is assessed below.

Plants for planting (trade) pathway

Various studies have reported the frequency with which Phytophthoras are associated with plants in nurseries and how such infected plants then act as a pathway to introduce these pathogens into natural, semi-natural and horticultural ecosystems (e.g., Jung et al. 2016). In the context of *P. pluvialis* the most likely pathway for introduction is on live conifer plants, particularly Douglas fir and radiata pine, although potentially other Pinus species could be hosts. In Great Britain, however, the Plant Health (Phytosanitary Conditions) (Amendment) (EU Exit) Regulations 2020, Annex 6, Part A (1.) prohibits the import of conifer plant species (Abies, Cedrus, Chamaecyparis, Juniperus, Larix, Picea, Pinus Pseudotsuga and Tsuga) other than as seeds, from all third countries other than EU member states and other specified European countries⁹. Identical requirements apply in Northern Ireland, under Commission Implementing Regulation (EU) 2019/2072 (Annex VI, item (1). Although the introduction of *P. pluvialis* into the UK probably pre-dates the most recent regulation, prohibition of conifer species from third countries has been in place for many years in the form of much earlier legislation applied to the Import and Export of Trees, Wood and Bark (Health) (Great Britain) Order 1980 and the Tree Pests (Great Britain) Order 1980. On this basis, entry via traded conifer plants for planting from North America (Canada, USA Pacific Northwest) and New Zealand where *P. pluvialis* is known to occur, is very unlikely. Despite this, there are uncertainties associated with this pathway

⁸ <u>https://forestry.gov.scot/sustainable-forestry/tree-health/tree-pests-and-diseases/phytophthora-pluvialis</u>

⁹ Specified European countries and areas are Albania, Andorra, Armenia, Azerbaijan, Belarus, Bosnia and Herzegovina, Canary Islands, Faroe Islands, Georgia, Iceland, Liechtenstein, Moldova, Monaco, Montenegro, North Macedonia, Norway, specific parts of Russia, San Marino, Serbia, Switzerland, Turkey and Ukraine.

such as the lack of information on the wider host range of *P. pluvialis*, especially in relation to broadleaf hosts. Additionally, although Tabima et al. (2021) suggest that *P. pluvialis* is likely to have originated in Oregon's coastal Douglas fir forests of the Pacific Northwest (PNW), with at least one introduction event from the PNW allowing it to establish in New Zealand, they also comment that "the potential exists for undiscovered populations of *P. pluvialis*" beyond the PNW and New Zealand.

Plants for planting



Overall, therefore, the plants for planting pathway is rated as **very unlikely** based on current and earlier import regulations applied to conifers although with **medium confidence** because of uncertainties about host range, particularly broadleaf species, and the wider geographical distribution of *P. pluvialis*.

Plants not for planting (non-trade) pathway

Due to the mild climate of southwest England and indeed much of western Britain, this region is rich in gardens open to the public which specialise in unusual or novel specimen ornamental shrubs and trees collected from many parts of the world¹⁰. Such plant collecting activities undertaken by both professionals and amateurs could potentially be a non-trade plant pathway for entry. Until 5-10 years ago, the biosecurity risks posed by these activities were rarely considered and collected plants brought in by this pathway were seldom quarantined or monitored after planting out to mitigate any risks of accidental pest introduction (Webber 2010). However, in comparison to the trade plant pathway, this non-trade plant pathway is relatively minor and sporadic, although likely to encompass a wide range of plant genera some of which be currently unknown hosts or non-traded hosts of *P. pluvialis* (e.g. tanoak). Considering the size of this pathway and the lack of awareness about associated biosecurity risks, it is rated as **unlikely** although with **low confidence** due to lack of data about plant numbers and uncertainty about the host range and geographical distribution of *P. pluvialis*.

Plants not for planting



¹⁰ Examples of plant collection gardens can be found at https://www.nationaltrust.org.uk/lists/top-gardens-to-visit-in-the-south-west



Timber/Wood pathway

Due to the export market for radiata pine grown in New Zealand, Hood et al. (2014) explored in depth the likelihood of movement of *P. pluvialis* on pine logs in the form of viable spores on bark surfaces or as colonised bark or sapwood. They concluded, based on loss of spore viability when tested as inoculum directly or in infected pine needles, coupled with the inability of *P. pluvialis* to colonise bark or sapwood of radiata pine, that neither logs or sawn timber were a likely pathway for the transport of the pathogen. Similar studies have not been undertaken with needles of Douglas fir, but stem lesion/canker formation has not been seen for this host in either New Zealand or the USA and spore viability is likely to be the same (Dick et al. 2014; Hansen et al. 2015).

Furthermore, any conifer wood or wood packaging material (WPM) to be imported into GB or Northern Ireland from EU or third countries including the USA and New Zealand must meet at least one of three phytosanitary requirements: (i) to be bark-free, (ii) to come from an area free from specified bark beetle pests or (iii) to be heat treated (56°C for 30 minutes). Hood et al. (2014) found that *P. pluvialis* oospores might not be viable after exposure for 2 hr at 35°C, suggesting higher temperatures (56°C for 30 minutes) would compound the effect.

Timber



The likelihood of entry via logs/wood/WPM is therefore assessed as **very unlikely** and with **high confidence** due to the poor ability of the pest to survive as a hitchhiker (possibly in the form of infected needles/needle fragments) on the exterior of wood, the lack of stem lesions on radiata pine or Douglas fir which infer that the pathogen is not present in bark or sapwood, and the phytosanitary treatments required for imported conifer wood or WPM. Stem cankers have only been recorded on tanoak on rare occasions (Reeser et al. 2013) and it is not a timber species, so again it is **very unlikely** that timber from this host would act as a pathway.

Cut Foliage pathway

Typically, foliage is collected from the wild for ornamental uses, it often comes from conifers, and is not subject to any specific cultural or treatment practices other than perhaps drying. As already described in Section 8, *P. pluvialis* has been found to produce abundant sporangia on infected needles of radiata pine (Gómez-Gallego et al. 2019a;) and by extension on infected needles of Douglas fir (Gómez-Gallego et al. 2019b). Therefore,

dried foliage or cut branches from susceptible hosts may contain viable pathogen structures, especially if formed within infected tissues, although drying (especially if using heat) is likely to reduce pathogen viability in or on the plant tissue. Material may also be dyed, bleached, or otherwise impregnated which may also reduce pathogen prevalence.

However, just as with plants for planting, the import of conifer foliage from all third countries is prohibited under GB Plant Health Regulations (see above). Additionally, oospores (likely to be the most resilient spore structures of *P. pluvialis*) have rarely been observed in infected needles of radiata pine (Hood et al. 2014; Williams & Hansen 2018), suggesting that survival by *P. pluvialis* during transport on this pathway is likely to be short-lived, especially if foliage is dried or subject to other treatments.

Therefore, the likelihood of entry via cut foliage is assessed as **very unlikely** although as cut foliage might also include hosts other than the currently known hosts, or Douglas fir for which there is a lack of information about the presence of oospores in infected needles, **medium confidence** is placed on this pathway rating.

Cut foliage



Seed pathway

In general, *Phytophthora* species are not considered to be seed borne pathogens. However, some examples exist: *Phytophthora cactorum* can be carried on beechnuts of *Fagus* (Prochazkova & Jancarek 1991), and a recent study demonstrated that seed infection by *Phytophthora gemini* of a common seagrass (*Zostera marina*) occurred frequently (Glovers et al. 2016). Even if seed is not infected directly, fruits or cones may become contaminated with soil, or be mixed with debris such as needles or other plant host material that contains pathogen propagules. Particularly with pathogens that infect needles, such as *P. pluvialis*, there is potential for conifer seed associated with infected debris to act as a pathway. Although plant health regulations prohibit the import of plants of conifer species (see above), seeds from North America and elsewhere are exempt from this measure. Since 1920, hundreds of kilos of seed collected from Douglas fir, western hemlock and radiata pine in North America have been imported into the UK (Anon 1965). More recently, data on forestry imports of conifer seed (Table 4) suggests that Douglas fir seed is still regularly imported into the UK from North America, so this pathway remains for this species at least.

Table 4: Quantity (in kg) of conifer seed imported into the UK from the USA and Canada since 2004(for years not shown in table no seed imports were recorded) (Forestry Commission Plant Health(2022)

	2004	2005	2008	2009	2010	2012	2014	2015	2016	2017	2018	2019	2020
--	------	------	------	------	------	------	------	------	------	------	------	------	------

Douglas fir	5	18	186	67	72	18	133	16	23	67	6	17	29
Western hemlock	0	0	0	0	0	0	0	0	0	0	0	0.5	0
Radiata pine	0	0	0	0	0	0	0	0	0	0	0	0	0

However, if few or no long-lived spore stages, such as oospores, are produced in needles by *P. pluvialis* (Hood et al. 2014; Williams & Hansen 2018) this would limit the ability to persist in contaminated soil/needle debris associated with seeds. For these reasons this pathway is assessed as **unlikely** but with **low confidence** in the rating due to lack of data of oospore formation in infected foliage of Douglas fir and the extent of debris contamination in imported seed lots.

Seed



Machinery and soil pathway

In speculating about the transfer of *P. pluvialis* from the coast range forests of Oregon to New Zealand, Tabima et al. (2021) suggest it could have occurred through various modes of introduction from infected Douglas fir needles to infested soil, due to lax biosecurity practices. In this context, Brar et al. (2017) mention that imported machinery could have been a possible pathway for *P. pluvialis*, as prior to 2001 the risks associated with used forestry machinery were not recognised in New Zealand (or probably elsewhere). As already described in Section 8, P. pluvialis has been found to produce abundant sporangia on infected needles of radiata pine (Gomez-Gallego et al. 2019a) and by extension on infected needles of Douglas fir (Gomez-Gallego et al. 2019a). As these needles die and fall to the forest floor often forming a thick carpet of prematurely shed needles, it creates potential for the pathogen to be harboured in soil and for contaminated soil to act as a pathway. However, as already indicated above, P. pluvialis may not persist for long in viable form in fallen needles and soil due to the lack of long-lived spore stages (Hood et al. 2014; Williams & Hansen 2018). In addition, importation of soil from third countries such as the USA and New Zealand is already prohibited under the Plant Health (Phytosanitary Conditions) (Amendment) (EU Exit) Regulations 2020 for Great Britain, and Commission Implementing Regulation (EU) 2019/2072 for Northern Ireland. However, machinery and vehicles which have been operated for forestry purposes can be imported providing they are accompanied by a phytosanitary certificate, and this should ensure that any risk is mitigated. Based on current regulation, this pathway is assessed as very unlikely, but with **medium confidence** due to the lack of data on how frequently used forestry machinery is imported into the UK from countries where *P. pluvialis* is present and levels

of compliance to make sure there is no associated soil/forestry debris with such machinery.



10. If the pest needs a vector, is it present in the UK/PRA area?

Phytophthora pluvialis does not require a vector for dispersal.

11. How likely is the pest to establish outdoors or under protection in the UK/PRA area?

Under protection

Establishment under protection is assessed as **very unlikely** with **high confidence**, because hosts of *P. pluvialis* are not grown under protection for their life span.

Under Protection	Very 🖌	Unlikely	Moderately Likely	Likely	Very 🗌 likely
Confidence	High Confidence	Medium Confidence	Low Confidence		

Outdoors

Host susceptibility and distribution influence the likelihood of *P. pluvialis* establishment in the wider environment; with climatic factors, such as temperature, moisture and needle wetness also playing a part in how likely outbreaks are to establish and the intensity at which they occur. Suitable hosts are also present in the UK (section 7; Figure 2).

In culture, *P. pluvialis* grows readily at between 15-20°C, with no growth above 25°C (Reeser et al. 2013), which is consistent with a lack of detections of *P. pluvialis* above a mean maximum temperature of 21.6°C in New Zealand (Fraser et al. 2020). Rainfall and relative humidity (RH) have also been highlighted as important drivers of RNC disease development and sporulation on radiata pine by Fraser et al. (2020). They found that sporulation and infection take place in the coolest, wettest part of the year (mid-winter through to mid-spring) peaking in June-August in New Zealand. At this time of year rainfall and RH are at their highest and average air temperatures at their lowest (Table 5; Scion 2022).

(Scion 2022)								
Month	Average Air Temp. (°C)	Average RH (%)	Average Total Rainfall (mm)					
Jan	17.01	79.33	93.9					
Feb	16.70	82.18	116.5					
Mar	14.72	86.72	201.7					
Apr	12.32	86.38	216.5					
May	10.49	84.82	180.4					
Jun	8.41	88.07	286.7					
Jul	7.24	86.94	251.9					
Aug	7.66	85.87	151.9					
Sep	8.95	83.59	292.5					
Oct	11.35	80.23	141.2					
Nov	13.31	79.88	132.2					
Dec	14.64	81.44	174.5					
Grand Average	11.89	83.79	186.7					

* Means for selected sites higher elevation coastal sites on the east coast of the North Island

Mild, wet winters, particularly those that are not too cold, are most likely to encourage the disease to establish, and in New Zealand it has been found that the most consistent disease expression is on higher elevation coastal sites (greater than 500m such as plateaus or ridge tops) on the east coast of the North Island with favourable relative humidity and rainfall (Scion 2022). The experience in the PNW with disease expression of *P. pluvialis* on Douglas fir is that it operates under similar climatic factors, except that the colder winters of the PNW tend to limit infections, as does the lower relative humidity in spring (Gómez-Gallego et al. 2019b) so overall levels of disease tend to be much lower compared with New Zealand and are also consistent with a native pathogen co-evolved with a native host.

In New Zealand and the cascades region of Oregon where *P. pluvialis* is active, the climate is probably very similar to western Britain. Indeed, many of the key commercial forestry conifers grown in Britain (Sitka spruce - *Picea sitchensis*, lodgepole pine - *Pinus contorta*, Douglas fir and western hemlock) have been chosen from the PNW because the climate similarities with Britain allow these timber producing species to thrive. Analysis of daily weather variables (temperature and rainfall) at a spatial resolution of 1km and using data available from 2001-2020, has visualised the parts of the UK where the risk of *P. pluvialis* establishment is likely to be greatest (Figure 3). The risk map shown below emphasises the suitability of climatic conditions throughout western Britain for *P. pluvialis* (Fera 2022). Additionally, the known hosts of *P. pluvialis* (Douglas fir, western hemlock and radiata pine) are most numerous in southwest England and Wales (Figure 2), where some of the most favourable climatic conditions occur. Between September 2021 and

March 2022, findings of *P. pluvialis* in the UK were made almost exclusively in western Britain, with locations ranging from the far north of Western Scotland (Ross-shire) down to southwest England (Cornwall). In addition, a single finding was made in Surrey where the climate is likely to less conducive, suggesting that establishment can occur even beyond the areas predicted to be most favourable for *P. pluvialis*.



Figure 3: Risk Map based on number of rainy days and temperature indicating areas likely to be most suitable for the establishment of *P. pluvialis.*

Overall, this confirms that a suitable climate for the pathogen is present through much of the PRA area. Similar climate conditions in the North Island of New Zealand, the PNW and particularly the western side of the UK, already favour other introduced aerial *Phytophthora* species such as *P. ramorum* (UK and PNW) and *P. kernoviae* (UK and NZ) and have allowed them to establish. Therefore, environmental conditions throughout at least western UK are predicted to be favourable to *P. pluvialis* so establishment is very likely and indeed has occurred already at some locations. On that basis the likelihood of establishment in the UK/PRA area is assessed as **very likely** with **high confidence**.



12. How quickly could the pest spread in the UK/PRA area?

Speed of spread will be influenced by several factors: the number of introductions and their spatial distribution, the number and distribution of suitable host plants, behaviour of the pest, and fluctuations in environmental conditions that influence pest behaviour and dispersal.

Natural spread

Although the biology of *P. pluvialis* is not fully understood, it is known that natural spread occurs via partially caducous sporangia that are produced on infected needles in the tree canopy (Dick et al. 2014; Williams & Hansen 2018). These easily detached sporangia are adapted for aerial dispersal in rain splash and fog. In New Zealand there has been no systematic work on the dispersal distances of sporangia, but most *P. pluvialis* spread is probably over short distances as illustrated by the localised disease expression within trees and forests. However, since RNC was first formally identified in New Zealand in 2008, findings of *P. pluvialis* have increased markedly (Graham et al. 2018). This, plus genotyping evidence that points to a single introduction followed by clonal spread (Tabima et al. 2021), suggests that spread by *P. pluvialis* has been moderately rapid over one to two decades. In addition, there are records of isolated trees and windbreaks that have become infected although some distance from another outbreak, indicating that longer distance aerial dispersal 'jumps' are also possible (Scion, 2022).

Apart from the arial dispersal of inoculum by fog and wind-driven rain, natural spread of *P. pluvialis* will also aided by movement of spores in water courses and run-off water, a

common occurrence with most Phytophthoras. Leaf baiting methods to detect *P. pluvialis* in water courses in Britain (see Section 7), have mainly been aimed at monitoring sporulation events, but inoculum has been detected at least 500 m downstream from areas with heavily diseased trees (Ana Pérez-Sierra, unpublished data). However, dilution effects as inoculum is moved away from disease foci in water courses probably limits pathogen spread through this process to relatively short distances (e.g. a few hundred metres), especially as the spore stages (sporangia and zoospores) are likely to be relatively short-lived and ephemeral.



On that basis, spread is considered likely to occur at a **moderate pace**, but this will be influenced by climate, the range and spatial distribution of hosts species, and possibly the influence of water courses around disease foci. Due to the lack of information on aerial dispersal distances and water course dispersal, confidence in this rating is placed at **medium**.

Spread via trade

A major pathway by which *P. pluvialis* is likely to spread (by analogy with other *Phytophthora* spp.) is on 'plants for planting' of known natural hosts but other host material such as timber and cut foliage may also sustain spread.

As only limited information is available on the full host range, there is also potential for spread on other species of *Pinus* as well as non-*Pinus* hosts. Currently there is no evidence to suggest *P. pluvialis* is circulating in the plant trade, despite the scattered outbreak findings in the UK (Section 6 – Summary). However, movement of infected plants in trade would allow long-distance jumps and new disease foci to be initiated so spread could occur **quickly** or even very quickly. Again, due to lack of information on the extent of the host range, and the pathogen source of the known outbreaks in the UK, there is **medium confidence** in this rating.

Plants for planting



Hood et al. (2014) assessed the likelihood of movement of *P. pluvialis* on radiata pine logs in the export trade from New Zealand and concluded the risk was minimal, but the

assessment was based on infected needles as the only source of inoculum and the inability of *P. pluvialis* to colonise bark or sapwood of radiata pine. This contrasts with the disease caused by *P. pluvialis* in Britain, in which not only the needles of western hemlock and Douglas fir are infected, but also bark in the form of cankers on twigs, branches and main stems of both host species (Pérez-Sierra et al. 2022a, b). The cankers which form on the main stem of both hosts can be extensive, with P. pluvialis invading phloem and cambial tissues and penetrating the sapwood. Initial assessments suggest sapwood penetration is superficial and limited to a few millimetres (J.F. Webber, unpublished data), but it potentially provides another opportunity for spread. The association with logs/timber may be (1) as a hitchhiker (possibly in the form of infected needles/needle fragments) on the exterior of logs although if the fragments are oospore-free then viability of *P. pluvialis* is likely to be short-lived (see Section 8), or (2) in infected phloem and sapwood underlying stem cankers. However, the ability of *P. pluvialis* to sporulate in canker tissues (and if so, what spore types are produced) is unknown and if limited/absent will reduce the potential for spread to new hosts. A suitable environment will also be required to encourage sporulation. Spread via timber or logs will therefore probably occur slowly but there is low confidence in this assessment, due to uncertainty in relation to the frequency, quantity, and type of sporulation.

Timber/logs



Foliage for ornamental uses (e.g. wreaths, flower arrangements) can be traded and may contain viable pathogen structures within infected tissues of susceptible hosts, although treatment of the foliage such as drying, dying, or bleaching is likely to reduce pathogen viability. After ornamental use, however, cut foliage is usually discarded into general waste which would limit potential for pathogen spread to susceptible hosts. On that basis, spread through traded foliage is considered likely to occur only **very slowly** with **high confidence** in this assessment.

Cut foliage



13. What is the pest's economic, environmental, and social impact within its existing distribution?

Douglas fir is considered one of the most important forest trees in Oregon (USA) and the second most valuable commercial conifer species for forestry in New Zealand; radiata pine is the most valuable conifer species in New Zealand. In comparisons of Douglas fir (New Zealand vs USA) evaluations indicate that *P. pluvialis* is much more abundant in New Zealand but causes much less conspicuous damage in the PNW (Gómez-Gallego et al. 2019b).

Reeser et al. (2015) suggest that P. pluvialis is "part of the native forest mycota of western Oregon", and its association with lower canopy needle loss in dense stands located in humid areas may be part of natural turnover, although P. pluvialis may also reduce the success of natural regeneration in dense Douglas fir stands. In New Zealand, however, the impact on plantation radiata pine and Douglas fir is of concern to forest growers in disease prone areas due to losses in increment growth that follows serious episodes of needle cast. Timing of disease expression and severity can also differ markedly between regions and years due to the influence of climate which makes guantification of increment losses associated with RNC difficult (Ganley et al. 2014). In an area where RNC was severe, the annual incremental growth of radiata pine decreased by ~35% in the following year, but growth losses do not appear to persist unless there is repeated defoliation. Williams & Hansen (2018) also comment that in some areas or years the incidence of RNC is so low that it is unlikely to result in any significant reductions in tree growth. Tree mortality due to P. pluvialis infection has never been observed in either Oregon or New Zealand on any host, but it cannot be ruled out that it may predispose trees to attack by secondary pests and pathogens (Ganley et al. 2014).

Based largely on the NZ experience with RNC where damage can be significant but intermittent, the impacts (economic/environmental/social) are rated as **medium** but with **low confidence**. There are major difficulties in rating the impact of a likely native pathogen co-evolved with a native host (Douglas fir) for which impacts are minor in the PNW, in combination with the same host and pathogen, but in the latter case where the pathogen has been introduced into a highly managed environment of plantation grown exotic tree species with differing environmental and social values (New Zealand). There is also a lack of quantitative data from both the PNW and NZ to make the assessment.



14. What is the pest's potential to cause economic, environmental, and social impacts in the UK/PRA area?

None of the known hosts of *P. pluvialis* are British native species, and the three most significantly affected (radiata pine, Douglas fir and western hemlock), are non-native species that originate from North America. The resource they provide is estimated at approximately 68,000 ha of forest cover in Britain, with an annual value of over £80 million, set within the context of a total of 1,308,000 ha of conifer cover in Britain. Only Douglas fir is considered a major forestry species of the known host species (see Section 7). Collectively, however, all pine species grown in Britain make up a much larger proportion of the plantation stock and if Scots or lodgepole pine proved to be susceptible, this would put a much larger number of trees and forest cover at risk (see Table 6 below).

Table 6: Areas (000 ha) and value of Potential *Phytophthora pluvialis* hosts in Great Britain shown by country¹¹ (Defra, unpublished data 2022)

Principal Species	England	Wales	Scotland	GB	Annual value of GB Woodlands (ONS GB Value)
Scots pine (Pinus sylvestris)	57	4	138	199	£237,052,017
Corsican pine (Pinus laricicola)	36	2	3	40*	£47,928,876
Lodgepole pine (Pinus contorta)	6	4	75	85*	£101,373,972
Douglas fir (<i>Pseudotsuga menziesii</i>)	25	9	25	59	£70,361,030
Western hemlock	6	1	2	9	£10,811,840
Other pines	28	4	31	63	£74,887,527
Total annual value					£542,415,262

* Totals may reflect rounding

Potential economic impacts in the PRA area

Economic impacts would not only depend on the number of tree species that might be susceptible to *P. pluvialis* infection, but also how severe disease symptoms could be and what parts of trees are affected. In the PNW and New Zealand, symptoms are limited to foliage, so hosts affected by severe episodes of *P. pluvialis* infection have temporarily reduced annual growth increment but apparently recover and are not killed by the pathogen. In contrast, *P. pluvialis* infection of western hemlock in the UK appears much more damaging, causing dieback and some mortality (unquantified), with cankers developing on the main tree stem that could affect the timber value at harvest. Symptoms

¹¹ Forestry Statistics 2021: <u>https://www.forestresearch.gov.uk/tools-and-resources/statistics/data-downloads/</u>

are less damaging on Douglas fir, but growth and therefore productivity is likely to be affected and stem cankers may again affect timber values. Currently *P. pluvialis* is being treated as a regulated pest, and tree felling is required with associated costs. The potential for timber salvage from the infected trees once felled is under investigation and will depend on the extent of sapwood penetration by *P. pluvialis* if infected wood has to be removed from the supply chain.

On that basis, economic impacts have the potential to be **large**, particularly if new/additional host species are found to be susceptible and become diseased. Annual value of conifer species at risk is estimated at over £500 million (Table 6), but uncertainty about the host range, the potential of affected trees to recover from infection episodes (some evidence is emerging to suggest this is happening) and whether it will be a continuing requirement to fell affected trees (plus any trees in a specified buffer zone) all affect the rating given for economic impacts. This is reflected in the **low confidence** rating applied to the assessment.



Potential environmental impacts in the PRA area

Environmental impacts include a range of ecosystem services. Thus, removal of large numbers of trees which might be required to deal with outbreaks could result in changes to soil (NH₄, NO₂ and cations), and on some sites soil erosion, changes in water quality and a decline in dissolved organic carbon. Carbon losses have been estimated to average around £70 per ha but could be as high as £490 per ha (2003 values from Willis et al. 2003). Changes to biodiversity are likely to include a loss of canopy resources and a possible short-term increase in saproxylic invertebrates (likely followed by a severe population decline due to a lack of suitable breeding material), although all three of the known conifer hosts are not native and endemic to Britain, so may have limited environmental value. There may, however, be an increase in ground flora (introduction of more light) particularly in hemlock stands which create dense understorey shade (Harmer et al. 2011). An average monetary value estimate for biodiversity loss is £146 per ha when estimated from Willis et al. (2003). More recent data on quantified environmental costs of the loss of mature trees are not available. Where conifer sites that are felled are scheduled for restoration under PAWS (returning Plantation sites to species mixes of Ancient Woodland Sites) with an emphasis on native broadleaf and conifer species in the mixed plantings, the result could be much greater biodiversity.

Environ -				
mental	Small	Medium 🗸	Large	
Impacts				



Environmental impacts are assessed as **medium**, but main tree species affected are not natives and it is uncertain if the requirement to fell affected trees plus those in a buffer zone will continue, so a **low confidence** rating applies to the assessment.

Potential social impacts in the PRA area

It is likely that there would be negative social responses to damage caused by *P. pluvialis* plus the impacts of widespread tree felling, due to the change in visual quality of affected forests especially if they are in areas used for amenity and recreational visits, although the latter would only apply to a limited proportion of affected trees. Clearing affected, potentially dying, trees for safety would also require priority action if they were in close proximity to recreational areas.

Overall social impacts are therefore judged as likely to be **medium** but with a **low confidence** due to the lack of data on the likely extent of damage and social analysis.



15. What is the pest's potential as a vector of plant pathogens?

Phytophthora pluvialis is a plant pathogen with no capacity to act as a vector of other pathogens.

16. What is the area endangered by the pest?

The current findings of *P. pluvialis* causing disease on hemlock and Douglas fir range from the north of west Scotland (Ross-shire), Wales and down to southwest England (Cornwall). This confirms that a suitable climate for the pathogen to establish and cause disease is present throughout much of the PRA area. *Phytophthora pluvialis* could potentially become established throughout the ranges of its known hosts in Britain (principally Douglas fir and western hemlock) which are found in woodlands, forest, parklands and gardens (National Forest Inventory <u>http://www.forestry.gov.uk/forestry/infd-8eyjwf</u>. Environmental conditions in the west of the UK are predicted to be most favourable to *P. pluvialis* for both establishment and spread (see Section 11).

Stage 3: Pest Risk Management

17. What are the risk management options for the UK/PRA area?

Exclusion

Recent findings indicate that *P. pluvialis* is already present in England, Scotland, and Wales at 35 separate locations (Section 6). It was classified by the UK Plant Health Risk Group, at least initially based on limited information about its UK distribution, as meeting the criteria to become a GB Quarantine Pest for regulatory purposes. This has enabled statutory action to be taken against findings and surveys are ongoing to assess how limited or widespread *P. pluvialis* is. Greater clarity about the extent of its distribution is required if Protected Zone status is to be considered for parts of the UK where the pest has not been found/reported (i.e. Northern Ireland) but exclusion through a Pest Free Area from the GB area does not currently apply.

The geographical distribution of the findings also suggests that *P. pluvialis* has been in Britain for some time, or that that have been multiple introductions although the level of risk identified around each pathway make this unlikely (see Section 9).

Eradication

Eradication is unlikely to be an effective option for *P. pluvialis* for various reasons. Eyre et al. (2022) set out some of the reasons for this which are:

- The number and widespread extent of the known outbreaks across Britain. The known extent of *P. pluvialis* already spans a significant area of southwest England and includes locations in Cumbria and several sites across Wales and western Scotland. In their analysis Pluess et al. (2012) found that the spatial extent of the initial infestation was significantly related to poor eradiation outcomes.
- The behaviour of the pathogen. Most *Phytophthora* species have highly successful survival strategies and tend to be supressed rather than eradicated by biocidal treatments (Jung et al. 2016).
- Eradication attempts with other *Phytophthora* introductions. There are no previous examples of non-native *Phytophthora* that have been found in the wider environment in the UK or possibly worldwide that have been successfully eradicated after establishment in the wider environment (Hansen 2015).
- The climatic limits of the pathogen. The presence of *P. pluvialis* in locations from southwest England to Northwest Scotland suggests that it is not at its climatic limits in Britain, especially as one of the outbreaks is in Surrey, England, and outside the most climatically suitable areas for establishment (see Figure 3). Eradication efforts are likely to be more challenging if the pathogen is not limited by climate.

Containment and controls

As an introduced pathogen in New Zealand, *P. pluvialis* has not been treated as a quarantine pest but efforts have been focussed on improving understanding of pathogen behaviour and management strategies. In New Zealand, it has become clear that disease outbreaks of RNC caused by *P. pluvialis* are cyclical and strongly weather dependent, reoccurring at different amplitudes every 2-3 years. However, these cycles are most apparent (and possibly most damaging) in new outbreak areas and less evident in areas where the pathogen is well-established (Dick et al. 2014; Gómez-Gallego et al. 2019b). Moreover, disease cycles are not associated with tree mortality and trees recover from even severe episodes of defoliation.

If eradication in the UK is not an option, then in the short-term, management options for consideration include:

- Containment of outbreak areas. Current evidence of the potential for natural spread by *P. pluvialis* is limited, but the process probably operates over limited distances (a few 100s of metres; see Section 12). Logs and timber from affected trees can carry the pathogen, but available evidence suggests the risk of spread from this material is limited and likely to be slow compared to spread to infected 'plants for planting'. Any risk of spread via harvested timber would be minimised through good biosecurity practices such as (1) jet washing machinery before it leaves affected areas to minimise the movement of contaminated/infected needles/debris through forestry operations, and (2) transfer any harvested material to already established, inspected bio-secure processors to prevent movement of infected bark/sapwood into the wood supply chain.
- Removal of affected hosts (Douglas fir and western hemlock). There is major uncertainty about the host range of *P. pluvialis*. Whilst more evidence is gathered on host range (e.g., susceptibility of all UK commercially grown pine species and some broadleaf species) and on dispersal potential under UK conditions, host removal could be focussed on the two known UK hosts at the worst affected locations. Felling and clearing of the affected overstorey and associated understorey would reduce build-up of sporulating material and therefore reduce the likelihood of natural spread through aerial dispersal of *P. pluvialis* sporangia/zoospores. Currently, little is known about sporulation potential on the foliage of different hosts (e.g. hemlock vs Douglas fir) and how this affects natural disease spread.
- Monitoring tree recovery and assessment of episodes of disease that arise in relation to known environmental triggers. This would extend under-standing of the impacts of *P. pluvialis* over time, for both significant outbreaks and only lightly affected outbreaks. There is emerging evidence that although *P. pluvialis* causes bark-killing cankers on branches and main stems of both western hemlock and Douglas fir, it is not unusual for canker development to halt and recovery to occur as callus occludes areas of cambial death caused by cankers (A. Pérez-Sierra and JF Webber, unpublished data). Recovery after disease episodes may be common, and on some affected sites canker

healing has been ongoing for 1-4 years. Additionally, despite confirmation of *P. pluvialis* at some locations, the pathogen is no longer active, only light symptoms are apparent and disease recovery is evident (A. Pérez-Sierra, unpublished data).

• Reviewing costs, impacts and benefits of host removal in relation to knowledge about host range and disease development.

Longer term options include:

- Trialling silvicultural treatments which change stand conditions, and by altering the microenvironment of infected trees reduce the likelihood of severe disease expression.
- Use of disease suppression compounds which reduce symptom expression, although such an approach would have to be consistent with certification and UKWAS. Phosphoric acid (phosphite) and copper oxychloride have both shown promise for the control of *P. pluvialis* in controlled inoculation experiments and within plantation systems although they have yet to be applied operationally within New Zealand forest systems. Aerial sprays of copper oxychloride are used routinely for other foliar pathogens (e.g., Dothistroma needle blight).
- Breeding for resistance as a long-term option for high value hosts. In New Zealand, field, and laboratory screening of radiata pine have shown that resistance to needle loss due to red needle cast is heritable and have identified resistance within current radiata pine breeding lines in use in New Zealand.

18. References

Abad G, Burgess T, Bienapfl JC, Redford AJ, Coffey M, Knight L. (2019). Molecular and morphological identification of *Phytophthora* species based on the types and other well-authenticated specimens. <u>https://idtools.org/id/phytophthora/index.php</u>

Anon. (1965). Seed identification numbers. Forestry Commission Research Branch Paper 29, 1-370.

Boevink PC, Birch PRJ, Turnbull D, Whisson SC. (2020). Devastating intimacy: the cell biology of plant–Phytophthora interactions. New Phytologist 228, 445–458 https/doi: 10.1111/nph.16650

Brar S, Tabima JF, McDougal RL, Dupont PY, Feau N, Hamelin RC, Panda P, LeBoldus JM, Grünwald NJ, Hansen EM, Bradshaw RE & Williams NM. (2017). Genetic diversity of *Phytophthora pluvialis*, a pathogen of conifers, in New Zealand and the west coast of the United States of America. Plant Pathology, 67, 1131–1139. https://doi.org/10.1111/ppa.12812 Dick MA, Williams NM, Bader M-K-F, Gardner JF & Bulman LS. (2014). Pathogenicity of *Phytophthora pluvialis* to *Pinus radiata* and its relation with red needle cast disease in New Zealand. Journal of Forestry Science 44(6) <u>https://doi.org/10.1186/s40490-014-0006-7</u>

Dungey HS, Williams NM, Low CB & Stovold GT. (2014). First evidence of genetic-based tolerance to red needle cast caused *by Phytophthora pluvialis* in radiata pine. New Zealand Journal of Forestry Science, 44. https://doi.org/10.1186/s40490-014-0028-1

EPPO (2022). Global Distribution Database – *Phytophthora pluvialis* (PHYTUV). <u>https://gd.eppo.int/taxon/PHYTUV/distribution Accessed 21/02/2022</u>

Erwin DC & Ribeiro OK. (1996). *Phytophthora* Diseases Worldwide. American Phytopathological Society Press, St Paul, MN.

Eyre D, Everatt M, Giltrap N & Lacey K. (2022). Comments on policy options document (17 Dec 2021) for *Phytophthora pluvialis* from risk managers in Defra's Risk and Horizon Scanning Team. Internal document.

Forestry Commission Plant Health (2022). Seed import data for *Pseudotsuga menziesii* and *Tsuga heterophylla*. Unpublished report. Or

Fraser S, Gomez-Gallego M, Gardner J, Bulman LS, Denman S & Williams NM. (2020). Impact of weather variables and season on sporulation of *Phytophthora pluvialis* and *Phytophthora kernoviae*. Forest Pathology, 50(n/a), e12588. https://doi.10.1111/efp.12588

Ganley RJ, Williams NM, Rolando CA, Hood IA, Dungey HS, Beets PN, Bulman LS. (2014). Management of red needle cast, caused by *Phytophthora pluvialis*, a new disease of radiata pine in New Zealand. New Zealand Plant Protection 67:48–53. Available from: <u>http://www.nzpps.org/nzpp_abstract.php?paper=670480</u>

Gardner J, Dobbie K, Fraser S. (2020). Shake it: how to collect Douglas fir needles for *Phytophthora* diagnostics. Forest Health News – Scion, No. 300, May 2020, ISSN 1175-9755

Govers LL, Man in 't Veld WA, Meffert JP, Bouma TJ, van Rijswick PCJ, Heusinkveld JHT, Orth RJ, van Katwijk MM, van der Heide T. (2016). Marine *Phytophthora* species can hamper conservation and restoration of vegetated coastal ecosystems. Proc. R. Soc. B 283: 20160812. <u>http://dx.doi.org/10.1098/rspb.2016.0812</u>

Gómez-Gallego M, Bader MK, Scott PM, Leuzinger S, Williams NM. (2017). *Phytophthora pluvialis* studies on Douglas-fir require Swiss needle cast suppression. Plant Disease 101(7),1259-1262. https://doi: 10.1094/PDIS-12-16-1738-RE

Gómez-Gallego M, Gommers R, Bader MK-F, Williams NM. (2019a). Modelling the key drivers of an aerial Phytophthora foliar disease epidemic, from the needles to the whole plant. PLoS ONE 14(5): e0216161

Gómez-Gallego M, LeBoldus JM, Bader M K-F, Hansen E, Donaldson L & Williams NM. (2019b). Contrasting the pathogen loads in co-existing populations of *Phytophthora pluvialis* and *Nothophaeocryptopus gaeumannii* in Douglas fir Plantations in New Zealand

and the Pacific Northwest United States. Phytopathology 109(11) <u>https://doi.org/10.1094/PHYTO-12-18-0479-R</u>

Hansen E. (2015). Phytophthora Species Emerging as Pathogens of Forest Trees. Current Forestry Reports 1. <u>https://doi:10.1007/s40725-015-0007-7</u>

Hansen EM, Reeser P, Sutton W, Gardner J, Williams N. (2015). First report of *Phytophthora pluvialis* causing needle loss and shoot dieback on Douglas-fir in Oregon and New Zealand. Plant Disease 99, 727.

Hansen E, Bourret T, Kanaskie A, LeBoldus L, Navarro S, Reeser P & Sutton W. (2017a). Current status of *Phytophthora pluvialis* in western North America. Proceedings of the 8th Meeting of the International Union of Forestry Research Organisations IUFRO Working Party S07-02-09 Phytophthora in Forests and Natural Ecosystems, Hanoi-Sapa Viet Nam March 2017 <u>http://forestphytophthoras.org/sites/default/files/proceedings/IUFRO-WP-7-02-09-Sapa-2017-abstracts-V2.pdf</u> page 24

Hansen EM, Reeser PW & Sutton W. (2017), Ecology and pathology of *Phytophthora* ITS clade 3 species in forests in western Oregon, USA, Mycologia 109(1), 100-114 http://doi: 10.1080/00275514.2016.1273622

Harmer R, Beauchamp K & Morgan G. (2011). Natural regeneration in western hemlock plantations on ancient woodland sites. Research Note 11: Forestry Commission, Edinburgh. <u>https://www.forestresearch.gov.uk/documents/295/FCRN011_ShIKdPO.pdf</u>

Hood I, Fraser S, Husheer S, Gardner J, Evanson T, Tieman G, Banham C & Wright L. (2022). Infection period of *Phytophthora pluvialis* and *Phytophthora kernoviae* in relation to weather variables in *Pinus radiata* forests in New Zealand. Submitted to journal.

Jung T et al. (2016). Widespread Phytophthora infestations in European nurseries put forest, semi-natural and horticultural ecosystems at high risk of Phytophthora diseases. Forest Pathology 46 (2):134-163. <u>https://doi.org/10.1111/efp.12239</u>

Pérez-Sierra A, Chitty R, Eacock A, Jones B, Biddle M, Crampton M, Lewis A, Olivieri L & Webber J. (2022). First report of *Phytophthora pluvialis* in Europe causing resinous cankers on western hemlock. New Disease Report, <u>https://doi.org/10.1002/ndr2.12064</u>

Pérez-Sierra A, Jones B, Biddle M & Webber J. (2022b). *Phytophthora pluvialis* – a new threat to forestry? Quarterly Journal of Forestry 116(2), 128-130.

Pluess T, Cannon R, Jarosik V, Pergl J, Pysek P & Bacher S. (2012). When are eradication campaigns successful? A test of common assumptions. Biological Invasions 14 (7):1365-1378. doi: 10.1007/s10530-011-0160-2.

Prochazkova Z & Jancarek V. (1991). Diseases in Czech-Slovak forest nurseries. In Proceedings of the first meeting of IUFRO working party S2.07-09 (Diseases and insects in forest nurseries, Pp. 37–50). (J.R. Sutherland and S. Glover, eds.). Inform. Rep. BC-X-331, Pacific For. Centre, For. Canada, Victoria, BC.

Reeser P, Sutton W & Hansen E. (2013). *Phytophthora pluvialis*, a new species from mixed tanoak-Douglas-fir forests of western Oregon, U.S.A. North American Fungi 8, 1–8. <u>https://doi.org/10.2509/naf2013.008.007</u>

Reeser P, Sutton W, Ganley B, Williams N, Hansen E. 2015. Phytophthora pluvialis. Forest Phytophthoras 5(1) <u>https://doi.10.5399/osu/fp.5.1.3745</u>

Scion (2022). *Phytophthora pluvialis* outbreak parameters in New Zealand. Unpublished report.

Scott PM, Taylor P & Williams N. (2019). Contrasting the infection and survival of *Phytophthora pluvialis* and *Phytophthora cinnamomi* in *Pinus radiata* roots. Australasian Plant Pathology <u>https://doi.10.1007/s13313-019-0619-7</u>

Webber JF. (2010). Pest risk analysis and invasion pathways for plant pathogens. New Zealand Journal of Forestry Science 40, 45-56.

Williams N & Hansen E. (2018). Red needle cast. In: Compendium of Conifer Diseases, Second Edition. (eds Hansen EM, Lewis KJ and Chastagner GA), pages 16-18. APS publications.

Willis K, Garrod G, Scarpa R, Powe N, Lovett A, Bateman I, Hanley N & Macmillan D. (2003). The social and environmental benefits of forests in Great Britain. Forestry Commission Report. 36pp. 2003. Edinburgh.

Name of Pest Risk Analysts(s)

Joan Webber, Forest Research

Joan.webber@forestresearch.gov.uk



© Crown copyright 2022

You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v.2. To view this licence visit <u>www.nationalarchives.gov.uk/doc/open-government-licence/version/2/</u> or email <u>PSI@nationalarchives.gov.uk</u>

This publication is available via the UK Plant Health Information portal <u>https://planthealthportal.defra.gov.uk/</u>

This PRA has been undertaken following IPPC International Standards for Phytosanitary Measures (ISPMs 2 and 11) and it provides technical evidence relating to the risk assessment and risk management of this pest.

Any enquiries regarding this publication should be sent to us at

The Chief Plant Health Officer

Department for Environment, Food and Rural Affairs

Room 11G32

Sand Hutton

York

YO41 1LZ

Email: plantpestsrisks@defra.gov.uk